

AP[®] Calculus AB Exam

SECTION I: Multiple Choice

2019

DO NOT OPEN THIS BOOKLET UNTIL YOU ARE TOLD TO DO SO.

At a Glance

Total Time

1 hour and 45 minutes

Number of Questions

45

Percent of Total Score

50%

Writing Instrument

Pencil required

Part A**Number of Questions**

30

Time

1 hour

Electronic Device

None allowed

Part B**Number of Questions**

15

Time

45 minutes

Electronic Device

Graphing calculator
required

Instructions

Section I of this exam contains 45 multiple-choice questions and 4 survey questions. For Part A, fill in only the circles for numbers 1 through 30 on the answer sheet. For Part B, fill in only the circles for numbers 76 through 90 on the answer sheet. Because Part A and Part B offer only four answer options for each question, do not mark the (E) answer circle for any question. The survey questions are numbers 91 through 94.

Indicate all of your answers to the multiple-choice questions on the answer sheet. No credit will be given for anything written in this exam booklet, but you may use the booklet for notes or scratch work. After you have decided which of the suggested answers is best, completely fill in the corresponding circle on the answer sheet. Give only one answer to each question. If you change an answer, be sure that the previous mark is erased completely. Here is a sample question and answer.

Sample Question Sample Answer

Chicago is a

(A) ☒ (B) ☐ (C) ☐ (D) ☐ (E) ☐

(A) state

(B) city

(C) country

(D) continent

Use your time effectively, working as quickly as you can without losing accuracy. Do not spend too much time on any one question. Go on to other questions and come back to the ones you have not answered if you have time. It is not expected that everyone will know the answers to all of the multiple-choice questions.

Your total score on the multiple-choice section is based only on the number of questions answered correctly. Points are not deducted for incorrect answers or unanswered questions.

Form I
Form Code 4PBP4-S

66

7. $\frac{d}{dx}(x^5 - 5^x) =$

(A) $\frac{x^6}{6} - \frac{5^x}{\ln 5}$ (B) $5x^4 - 5^x$ (C) $5x^4 - x \cdot 5^{x-1}$ (D) $5x^4 - (\ln 5)5^x$


8. $\lim_{x \rightarrow \infty} \frac{10 - 6x^2}{5 + 3e^x}$ is

(A) -2 (B) 0 (C) 2 (D) nonexistent

18. $\frac{d}{dx}(x^3 \sec(2x)) =$

- (A) $6x^2 \sec(2x) \tan(2x)$
- (B) $2x^3 \tan^2(2x) + 3x^2 \sec(2x)$
- (C) $x^3 \sec(2x) \tan(2x) + 3x^2 \sec(2x)$
- (D) $2x^3 \sec(2x) \tan(2x) + 3x^2 \sec(2x)$

-
19. A particle moves along the y-axis so that at time $t \geq 0$ its position is given by $y(t) = t^3 - 4t^2 + 4t + 3$. Which of the following statements describes the motion of the particle at time $t = 1$?
- (A) The particle is moving down the y-axis with decreasing velocity.
 - (B) The particle is moving down the y-axis with increasing velocity.
 - (C) The particle is moving up the y-axis with decreasing velocity.
 - (D) The particle is moving up the y-axis with increasing velocity.



28. A person stands 30 feet from point P and watches a balloon rise vertically from the point, as shown in the figure above. The balloon is rising at a constant rate of 2 feet per second. What is the rate of change, in radians per second, of angle θ at the instant when the balloon is 40 feet above point P ?

(A) $\frac{3}{100}$ (B) $\frac{3}{125}$ (C) $\frac{1}{12}$ (D) $\frac{5}{27}$

-
29. How many vertical asymptotes does the graph of $y = \frac{x - 2}{x^4 - 16}$ have?

(A) One (B) Two (C) Three (D) Four

30. For what value of b does the integral $\int_1^b x^2 dx$ equal $\lim_{n \rightarrow \infty} \sum_{k=1}^n \left(1 + \frac{2k}{n}\right)^2 \frac{2}{n}$?

- 23-

B**B****B****B****B****B****B****B****B****CALCULUS AB****SECTION I, Part B****Time—45 minutes****Number of questions—15****A GRAPHING CALCULATOR IS REQUIRED FOR SOME QUESTIONS ON THIS PART OF THE EXAM.**


Directions: Solve each of the following problems, using the available space for scratch work. After examining the form of the choices, decide which is the best of the choices given and fill in the corresponding circle on the answer sheet. No credit will be given for anything written in this exam booklet. Do not spend too much time on any one problem.

BE SURE YOU FILL IN THE CIRCLES ON THE ANSWER SHEET THAT CORRESPOND TO QUESTIONS NUMBERED 76–90.

YOU MAY NOT RETURN TO QUESTIONS NUMBERED 1–30.

In this exam:

- (1) The exact numerical value of the correct answer does not always appear among the choices given. When this happens, select from among the choices the number that best approximates the exact numerical value.
- (2) Unless otherwise specified, the domain of a function f is assumed to be the set of all real numbers x for which $f(x)$ is a real number.
- (3) The inverse of a trigonometric function f may be indicated using the inverse function notation f^{-1} or with the prefix “arc” (e.g., $\sin^{-1} x = \arcsin x$).

B**B****B****B****B****B****B****B****B**

76. The function f is continuous on the closed interval $[0, 5]$. The graph of f' , the derivative of f , is shown above. On which of the following intervals is f increasing?
- (A) $[0, 1]$ and $[2, 4]$
- (B) $[0, 1]$ and $[3, 5]$
- (C) $[0, 1]$ and $[4, 5]$ only
- (D) $[0, 2]$ and $[4, 5]$

B**B****B****B****B****B****B****B****B**

77. The height of an object at time $t \geq 1$ is given by $h(t) = t^2 - \frac{16}{t} + 15$. What is the velocity of the object at time $t = 3$?

- (A) 0.815 (B) 7.778 (C) 18.667 (D) 21.089

78. The function g is differentiable and satisfies $g(-1) = 4$ and $g'(-1) = 2$. What is the approximation of $g(-1.2)$ using the line tangent to the graph of g at $x = -1$?

- (A) 3.6 (B) 3.8 (C) 4.2 (D) 4.4

B**B****B****B****B****B****B****B****B**

79. Tara's heart rate during a workout is modeled by the differentiable function h , where $h(t)$ is measured in beats per minute and t is measured in minutes from the start of the workout. Which of the following expressions gives Tara's average heart rate from $t = 30$ to $t = 60$?

- (A) $\int_{30}^{60} h(t) \, dt$
(B) $\frac{1}{30} \int_{30}^{60} h(t) \, dt$
(C) $\frac{1}{30} \int_{30}^{60} h'(t) \, dt$
(D) $\frac{h'(30) + h'(60)}{2}$


-
80. Let g be the function with first derivative $g'(x) = \sqrt{x^3 + x}$ for $x > 0$. If $g(2) = -7$, what is the value of $g(5)$?

- (A) 4.402 (B) 11.402 (C) 13.899 (D) 20.899


B**B****B****B****B****B****B****B****B**

81. If f is a function that has a removable discontinuity at $x = 3$, which of the following could be the graph of f ?


(A)




(B)



(C)



(D)



B**B****B****B****B****B****B****B****B**

82. Let f be a continuous function such that $\int_0^{17} f(x) \, dx = 8$, $\int_{17}^{20} f(x) \, dx = -3$, and $\int_{13}^{20} f(x) \, dx = 7$. What is the value of $\int_0^{13} f(x) \, dx$?

(A) -2 (B) 4 (C) 12 (D) 18

$$f(x) = \begin{cases} k^3 + x & \text{for } x < 3 \\ \frac{16}{k^2 - x} & \text{for } x \geq 3 \end{cases}$$


83. Let f be the function defined above, where k is a positive constant. For what value of k , if any, is f continuous?
- (A) 2.081 (B) 2.646 (C) 8.550 (D) There is no such value of k .

B**B****B****B****B****B****B****B****B**

x	$f(x)$
-5	-9
0	1
2	5


84. The table above gives values of a continuous function f at selected values of x . Based on the information in the table, which of the following statements must be true?
- (A) f has at most one zero.
- (B) f has a relative maximum at $x = 2$.
- (C) There exists a value c , where $-5 < c < 2$, such that $f(c) = 4$.
- (D) There exists a value c , where $-5 < c < 2$, such that $f'(c) = 2$.
-

85. The function f is an antiderivative of the function g defined by $g(x) = 3 - \sqrt{x^2 + x + 4 \cos x}$. Which of the following is the x -coordinate of the location of a local maximum for the graph of $y = f(x)$?
- (A) -3.961 (B) -2.161 (C) 1.494 (D) 3.140


B**B****B****B****B****B****B****B****B**

86. The graph of $y = f(x)$ on the interval $0 < x < 5$ is shown above. Which of the following could be the graph of $y = f'(x)$?


(A)




(B)



(C)



(D)



B**B****B****B****B****B****B****B****B**

87. At time t , $0 < t < 2$, the velocity of a particle moving along the x -axis is given by $v(t) = t \sin(t^3)$. Let $t = b$ be the time at which the particle changes direction from moving left to moving right. What is the total distance traveled by the particle during the time interval $0 < t < b$?
- (A) 0.212 (B) 0.612 (C) 1.011 (D) 1.208

-
88. Let f be the function defined by $f(x) = \frac{1}{4}x^4 - \frac{2}{3}x^3 + \frac{1}{2}x^2 - \frac{1}{2}x$. For how many values of x in the open interval $(0, 1.565)$ is the instantaneous rate of change of f equal to the average rate of change of f on the closed interval $[0, 1.565]$?
- (A) Zero (B) One (C) Three (D) Four

B**B****B****B****B****B****B****B****B**

89. Let g be a twice-differentiable function with $g'(x) > 0$ and $g''(x) > 0$ for all real numbers x , such that $g(3) = 12$ and $g(5) = 18$. Which of 20, 21, and 22 are possible values for $g(6)$?
- (A) 21 only (B) 22 only (C) 20 and 21 only (D) 21 and 22 only

B**B****B****B****B****B****B****B****B**

x	2	3	4
$f(x)$	1	2	6
$f'(x)$	4	5	3

90. The table above gives values of the differentiable function f and its derivative at selected values of x . If g is the inverse function of f , which of the following is an equation of the line tangent to the graph of g at the point where $x = 2$?

(A) $y = -\frac{1}{5}(x - 2) + 3$

(B) $y = -\frac{1}{4}(x - 2) + 1$

(C) $y = \frac{1}{5}(x - 2) + 3$

(D) $y = 4(x - 2) + 1$

B**B****B****B****B****B****B****B****B**

END OF SECTION I

**IF YOU FINISH BEFORE TIME IS CALLED,
YOU MAY CHECK YOUR WORK ON PART B ONLY.**

DO NOT GO ON TO SECTION II UNTIL YOU ARE TOLD TO DO SO.

MAKE SURE YOU HAVE DONE THE FOLLOWING.

- **PLACED YOUR AP NUMBER LABEL ON YOUR ANSWER SHEET**
- **WRITTEN AND GRIDDED YOUR AP NUMBER CORRECTLY ON YOUR ANSWER SHEET**
- **TAKEN THE AP EXAM LABEL FROM THE FRONT OF THIS BOOKLET AND PLACED IT ON YOUR ANSWER SHEET**

AP[®] Calculus AB Exam

SECTION II: Free Response

2019

DO NOT OPEN THIS BOOKLET OR BREAK THE SEALS ON PART B UNTIL YOU ARE TOLD TO DO SO.

At a Glance

Total Time

1 hour and 30 minutes

Number of Questions

6

Percent of Total Score

50%

Writing Instrument

Either pencil or pen with black or dark blue ink

Weight

The questions are weighted equally, but the parts of a question are not necessarily given equal weight.

Part A

Number of Questions

2

Time

30 minutes

Electronic Device

Graphing calculator required

Percent of Section II Score

33.33%

Part B

Number of Questions

4

Time

1 hour

Electronic Device

None allowed

Percent of Section II Score

66.67%

IMPORTANT Identification Information

PLEASE PRINT WITH PEN:

1. First two letters of your last name

First letter of your first name

2. Date of birth

Month Day Year

3. Six-digit school code

4. Unless I check the box below, I grant the College Board the unlimited right to use, reproduce, and publish my free-response materials, both written and oral, for educational research and instructional purposes. My name and the name of my school will not be used in any way in connection with my free-response materials. I understand that I am free to mark "No" with no effect on my score or its reporting.

No, I do not grant the College Board these rights. ☐

Instructions

The questions for Section II are printed in this booklet. Do not break the seals on Part B until you are told to do so. Write your solution to each part of each question in the space provided. Write clearly and legibly. Cross out any errors you make; erased or crossed-out work will not be scored.

Manage your time carefully. During Part A, work only on the questions in Part A. You are permitted to use your calculator to solve an equation, find the derivative of a function at a point, or calculate the value of a definite integral. However, you must clearly indicate the setup of your question, namely the equation, function, or integral you are using. If you use other built-in features or programs, you must show the mathematical steps necessary to produce your results. During Part B, you may continue to work on the questions in Part A without the use of a calculator.

As you begin each part, you may wish to look over the questions before starting to work on them. It is not expected that everyone will be able to complete all parts of all questions.

- Show all of your work, even though a question may not explicitly remind you to do so. Clearly label any functions, graphs, tables, or other objects that you use. Justifications require that you give mathematical reasons, and that you verify the needed conditions under which relevant theorems, properties, definitions, or tests are applied. Your work will be scored on the correctness and completeness of your methods as well as your answers. Answers without supporting work will usually not receive credit.
- Your work must be expressed in standard mathematical notation rather than calculator syntax. For example, $\int_1^5 x^2 dx$ may not be written as `fnInt(X2, X, 1, 5)`.
- Unless otherwise specified, answers (numeric or algebraic) need not be simplified. If you use decimal approximations in calculations, your work will be scored on accuracy. Unless otherwise specified, your final answers should be accurate to three places after the decimal point.
- Unless otherwise specified, the domain of a function f is assumed to be the set of all real numbers x for which $f(x)$ is a real number.

Form I

Form Code 4BP4-S

66

CALCULUS AB
SECTION II, Part A
Time—30 minutes
Number of questions—2

A GRAPHING CALCULATOR IS REQUIRED FOR THESE QUESTIONS.

t (hours)	2	5	9	11	12
$L(t)$ (cars per hour)	15	40	24	68	18

1. The rate at which cars enter a parking lot is modeled by $E(t) = 30 + 5(t - 2)(t - 5)e^{-0.2t}$. The rate at which cars leave the parking lot is modeled by the differentiable function L . Selected values of $L(t)$ are given in the table above. Both $E(t)$ and $L(t)$ are measured in cars per hour, and time t is measured in hours after 5 A.M. ($t = 0$). Both functions are defined for $0 \leq t \leq 12$.

(a) What is the rate of change of $E(t)$ at time $t = 7$? Indicate units of measure.

- (b) How many cars enter the parking lot from time $t = 0$ to time $t = 12$? Give your answer to the nearest whole number.

Do not write beyond this border.

Do not write beyond this border.



- (c) Use a trapezoidal sum with the four subintervals indicated by the data in the table to approximate

$\int_2^{12} L(t) dt$. Using correct units, explain the meaning of $\int_2^{12} L(t) dt$ in the context of this problem.

-
- (d) For $0 \leq t < 6$, 5 dollars are collected from each car entering the parking lot. For $6 \leq t \leq 12$, 8 dollars are collected from each car entering the parking lot. How many dollars are collected from the cars entering the parking lot from time $t = 0$ to time $t = 12$? Give your answer to the nearest whole dollar.

Do not write beyond this border.

Do not write beyond this border.



2. The function f is defined by $f(x) = 3(1+x)^{0.5} \cos\left(\frac{\pi x}{6}\right)$ for $0 \leq x \leq 3$. The function g is continuous and decreasing for $0 \leq x \leq 3$ with $g(3) = 0$.

The figure above on the left shows the graphs of f and g and the regions R and S . R is the region bounded by the graph of g and the x - and y -axes. Region R has area 3.24125. S is the region bounded by the y -axis and the graphs of f and g .

The figure above on the right shows the graph of $y = (g(x))^2$ and the region T . T is the region bounded by the graph of $y = (g(x))^2$ and the x - and y -axes. Region T has area 5.32021.

- (a) Find the area of region S .

- (b) Find the volume of the solid generated when region S is revolved about the horizontal line $y = -3$.

-
- (c) Region S is the base of a solid. For this solid, each cross section perpendicular to the x -axis is a rectangle whose height is 7 times the length of its base in region S . Write, but do not evaluate, an integral expression for the volume of this solid.

Do not write beyond this border.

Do not write beyond this border.

END OF PART A

**IF YOU FINISH BEFORE TIME IS CALLED,
YOU MAY CHECK YOUR WORK ON PART A ONLY.
DO NOT GO ON TO PART B UNTIL YOU ARE TOLD TO DO SO.**

CALCULUS AB
SECTION II, Part B
Time—1 hour
Number of questions—4

NO CALCULATOR IS ALLOWED FOR THESE QUESTIONS.

DO NOT BREAK THE SEALS UNTIL YOU ARE TOLD TO DO SO.

NO CALCULATOR ALLOWED

$$f(x) = \begin{cases} \sqrt{9 - x^2} & \text{for } -3 \leq x \leq 0 \\ -x + 3 \cos\left(\frac{\pi x}{2}\right) & \text{for } 0 < x \leq 4 \end{cases}$$

3. Let f be the function defined above.

(a) Find the average rate of change of f on the interval $-3 \leq x \leq 4$.

(b) Write an equation for the line tangent to the graph of f at $x = 3$.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED


(c) Find the average value of f on the interval $-3 \leq x \leq 4$.

(d) Must there be a value of x at which $f(x)$ attains an absolute maximum on the closed interval $-3 \leq x \leq 4$? Justify your answer.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED

Graph of f

4. The continuous function f is defined for $-4 \leq x \leq 4$. The graph of f , shown above, consists of two line segments and portions of three parabolas. The graph has horizontal tangents at $x = -\frac{1}{2}$, $x = \frac{1}{2}$, and $x = \frac{5}{2}$. It is known that $f(x) = -x^2 + 5x - 4$ for $1 \leq x \leq 4$. The areas of regions A and B bounded by the graph of f and the x -axis are 3 and 5, respectively. Let g be the function defined by $g(x) = \int_{-4}^x f(t) dt$.
- (a) Find $g(0)$ and $g(4)$.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED

(b) Find the absolute minimum value of g on the closed interval $[-4, 4]$. Justify your answer.

(c) Find all intervals on which the graph of g is concave down. Give a reason for your answer.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED

t (hours)	0	1	2	3	4
$B(t)$ (miles per hour)	1	8	1.5	-5	11

5. Brandon and Chloe ride their bikes for 4 hours along a flat, straight road. Brandon's velocity, in miles per hour, at time t hours is given by a differentiable function B for $0 \leq t \leq 4$. Values of $B(t)$ for selected times t are given in the table above. Chloe's velocity, in miles per hour, at time t hours is given by the piecewise function C defined by

$$C(t) = \begin{cases} te^{4-t^2} & \text{for } 0 \leq t \leq 2 \\ 12 - 3t - t^2 & \text{for } 2 < t \leq 4. \end{cases}$$

- (a) How many miles did Chloe travel from time $t = 0$ to time $t = 2$?

-
- (b) At time $t = 3$, is Chloe's speed increasing or decreasing? Give a reason for your answer.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED

- (c) Is there a time t , for $0 \leq t \leq 4$, at which Brandon's acceleration is equal to 2.5 miles per hour per hour? Justify your answer.

-
- (d) Is there a time t , for $0 \leq t \leq 2$, at which Brandon's velocity is equal to Chloe's velocity? Justify your answer.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED

6. Consider the curve defined by $2x^2 + 3y^2 - 4xy = 36$.

(a) Show that $\frac{dy}{dx} = \frac{2y - 2x}{3y - 2x}$.

(b) Find the slope of the line tangent to the curve at each point on the curve where $x = 6$.

Do not write beyond this border.

Do not write beyond this border.

NO CALCULATOR ALLOWED

- (c) Find the positive value of x at which the curve has a vertical tangent line. Show the work that leads to your answer.

-
- (d) Let x and y be functions of time t that are related by the equation $2x^2 + 3y^2 - 4xy = 36$. At time $t = 1$, the value of x is 2, the value of y is -2 , and the value of $\frac{dy}{dt}$ is 4. Find the value of $\frac{dx}{dt}$ at time $t = 1$.

Do not write beyond this border.

Do not write beyond this border.

STOP
END OF EXAM

THE FOLLOWING INSTRUCTIONS APPLY TO THE COVERS OF THE SECTION II BOOKLET.

- **MAKE SURE YOU HAVE COMPLETED THE IDENTIFICATION INFORMATION AS REQUESTED ON THE FRONT AND BACK COVERS OF THE SECTION II BOOKLET.**
- **CHECK TO SEE THAT YOUR AP NUMBER LABEL APPEARS IN THE BOX ON THE FRONT COVER.**
- **MAKE SURE YOU HAVE USED THE SAME SET OF AP NUMBER LABELS ON ALL AP EXAMS YOU HAVE TAKEN THIS YEAR.**

Answer Key for AP Calculus AB
Practice Exam, Section I

Question 1: B	Question 76: D
Question 2: C	Question 77: B
Question 3: D	Question 78: A
Question 4: D	Question 79: B
Question 5: C	Question 80: C
Question 6: D	Question 81: C
Question 7: D	Question 82: A
Question 8: B	Question 83: A
Question 9: B	Question 84: C
Question 10: A	Question 85: D
Question 11: A	Question 86: A
Question 12: C	Question 87: C
Question 13: A	Question 88: C
Question 14: D	Question 89: B
Question 15: A	Question 90: C
Question 16: B	
Question 17: B	
Question 18: D	
Question 19: A	
Question 20: C	
Question 21: D	
Question 22: D	
Question 23: B	
Question 24: A	
Question 25: A	
Question 26: B	
Question 27: C	
Question 28: B	
Question 29: A	
Question 30: B	

Multiple-Choice Section for Calculus AB
2019 Course Framework Alignment and Rationales

Question 1

Skill	Learning Objective	Topic
1.E	FUN-6.C	Finding Antiderivatives and Indefinite Integrals - Basic Rules and Notation
(A)	Incorrect. This is the derivative of $\frac{x^2}{4}$, not the antiderivative.	
(B)	Correct. By the power rule for antiderivatives, the antiderivative of x^n is $\frac{x^{n+1}}{n+1}$ for $n \neq -1$. Therefore, $\int \frac{x^2}{4} dx = \frac{1}{4} \int x^2 dx = \frac{1}{4} \cdot \frac{x^3}{3} + C = \frac{x^3}{12} + C.$	
(C)	Incorrect. This response would result if the power rule for antiderivatives was not applied correctly. The antiderivative of x^2 was taken to be x^3 rather than $\frac{x^3}{3}$.	
(D)	Incorrect. This response would result if the power rule for antiderivatives was not applied correctly. The antiderivative of x^2 was taken to be $3x^3$ rather than $\frac{x^3}{3}$.	

Question 2

Skill	Learning Objective	Topic
1.D	CHA-2.C	Defining the Derivative of a Function and Using Derivative Notation
(A)	Incorrect. This response would result if the derivative of $\cos x$ was taken to be $\sin x$ rather than $-\sin x$. The slope at the point $\left(\frac{\pi}{2}, 0\right)$ was therefore taken to be 1. In addition, an equation of a line through the point (x_0, y_0) was written as $y = m(x + x_0) + y_0$ instead of $y = m(x - x_0) + y_0$, leading to the response $y = +1\left(x + \frac{\pi}{2}\right) + 0 = x + \frac{\pi}{2}.$	
(B)	Incorrect. This response would result if the derivative of $\cos x$ was taken to be $\sin x$ rather than $-\sin x$. The slope at the point $\left(\frac{\pi}{2}, 0\right)$ was therefore taken to be 1, giving an equation of the tangent line as $y = +1\left(x - \frac{\pi}{2}\right) + 0 = x - \frac{\pi}{2}.$	
(C)	Correct. The slope of the tangent line is the value of the derivative at $x = \frac{\pi}{2}$. $\frac{dy}{dx} = -\sin x \Rightarrow \frac{dy}{dx}\bigg _{x=\frac{\pi}{2}} = -\sin\left(\frac{\pi}{2}\right) = -1$ At $x = \frac{\pi}{2}$, $y = \cos\left(\frac{\pi}{2}\right) = 0$. An equation of the tangent line at the point $\left(\frac{\pi}{2}, 0\right)$ is therefore $y = -1\left(x - \frac{\pi}{2}\right) + 0 = -x + \frac{\pi}{2}.$	
(D)	Incorrect. This response might come from writing an equation of a line through the point (x_0, y_0) as $y = m(x + x_0) + y_0$ instead of $y = m(x - x_0) + y_0$, leading to $y = -1\left(x + \frac{\pi}{2}\right) + 0 = -x - \frac{\pi}{2}$. Alternately, an error might have been made in simplifying an equation of the tangent line. The slope of the tangent line at $x = \frac{\pi}{2}$ was correctly found. $\frac{dy}{dx} = -\sin x \Rightarrow \frac{dy}{dx}\bigg _{x=\frac{\pi}{2}} = -\sin\left(\frac{\pi}{2}\right) = -1$ An equation of the tangent line at the point $\left(\frac{\pi}{2}, 0\right)$ was written, as follows. $y = -1\left(x - \frac{\pi}{2}\right) + 0 = -x - \frac{\pi}{2}$	

Question 3

Skill	Learning Objective	Topic
1.E	FUN-3.C	The Chain Rule
(A)	Incorrect. This response might come from incorrectly applying the chain rule twice as $\frac{d}{dx}(f(g(x))) = f'(g'(x))$, as follows. $\frac{d}{dx}\left(2(\sin\sqrt{x})^2\right) = 2 \cdot \left(2 \cdot \frac{d}{dx}(\sin\sqrt{x})\right) = 2 \cdot 2 \cdot \cos\left(\frac{d}{dx}(\sqrt{x})\right) = 4\cos\left(\frac{1}{2\sqrt{x}}\right)$	
(B)	Incorrect. This response might come from correctly applying the chain rule the first time but not the second, as follows. $\frac{d}{dx}\left(2(\sin\sqrt{x})^2\right) = 2 \cdot 2(\sin\sqrt{x}) \cdot \frac{d}{dx}(\sin\sqrt{x}) = 2 \cdot 2(\sin\sqrt{x}) \cdot \cos\sqrt{x}$	
(C)	Incorrect. This response might come from using the chain rule only once, with the innermost “inside” function, $\sqrt{x}$, as follows. $\frac{d}{dx}\left(2(\sin\sqrt{x})^2\right) = 2 \cdot 2(\sin\sqrt{x}) \cdot \frac{d}{dx}(\sqrt{x}) = 2 \cdot 2(\sin\sqrt{x}) \cdot \frac{1}{2\sqrt{x}}$	
(D)	Correct. The chain rule must be used twice for this composition of three functions. $\begin{aligned} \frac{d}{dx}\left(2(\sin\sqrt{x})^2\right) &= 2 \cdot 2(\sin\sqrt{x}) \cdot \left(\frac{d}{dx}(\sin\sqrt{x})\right) \\ &= 2 \cdot 2(\sin\sqrt{x}) \cdot \left(\cos\sqrt{x} \cdot \frac{d}{dx}(\sqrt{x})\right) \\ &= 2 \cdot 2(\sin\sqrt{x}) \cdot \cos\sqrt{x} \cdot \frac{1}{2\sqrt{x}} \\ &= \frac{2\sin\sqrt{x}\cos\sqrt{x}}{\sqrt{x}} \end{aligned}$	

Question 4

Skill	Learning Objective	Topic
1.E	FUN-6.D	Integrating Using Substitution
(A)	Incorrect. This response would result if the factor 3 was mishandled during the substitution using $u = x^3 + 3x - 5$, and the expression $3x^2 + 3$ from the derivative was substituted back for u rather than the expression $x^3 + 3x - 5$, as follows. $u = x^3 + 3x - 5 \Rightarrow \frac{du}{dx} = 3x^2 + 3 = 3(x^2 + 1) \Rightarrow dx = \frac{3 du}{(x^2 + 1)}$ $\int \frac{x^2 + 1}{(x^3 + 3x - 5)^3} dx = \int \frac{1}{u^3} \cdot 3 du = 3 \cdot \left(-\frac{1}{2u^2} \right) + C = -\frac{3}{2} \cdot \frac{1}{(3x^2 + 3)^2} + C$	
(B)	Incorrect. Starting with the substitution $u = x^3 + 3x - 5$, $u = x^3 + 3x - 5 \Rightarrow \frac{du}{dx} = 3x^2 + 3 = 3(x^2 + 1) \Rightarrow dx = \frac{du}{3(x^2 + 1)}.$ Substituting for $x^3 + 3x - 5$ and for dx gives $\int \frac{x^2 + 1}{(x^3 + 3x - 5)^3} dx = \int \frac{1}{u^3} \cdot \frac{1}{3} du = \frac{1}{3} \cdot \left(-\frac{1}{2u^2} \right) + C.$ The expression $3x^2 + 3$ from the derivative might have been substituted back for u, however, rather than the expression $x^3 + 3x - 5$.	
(C)	Incorrect. This response would result if the factor 3 was mishandled during the substitution using $u = x^3 + 3x - 5$, as follows. $u = x^3 + 3x - 5 \Rightarrow \frac{du}{dx} = 3x^2 + 3 = 3(x^2 + 1) \Rightarrow dx = \frac{3du}{(x^2 + 1)}$ $\int \frac{x^2 + 1}{(x^3 + 3x - 5)^3} dx = \int \frac{1}{u^3} \cdot 3 du = 3 \cdot \left(-\frac{1}{2u^2} \right) + C = -\frac{3}{2} \cdot \frac{1}{(x^3 + 3x - 5)^2} + C$	
(D)	Correct. Starting with the substitution $u = x^3 + 3x - 5$, $u = x^3 + 3x - 5 \Rightarrow \frac{du}{dx} = 3x^2 + 3 = 3(x^2 + 1) \Rightarrow dx = \frac{du}{3(x^2 + 1)}.$ Substituting for $x^3 + 3x - 5$ and for dx gives $\int \frac{x^2 + 1}{(x^3 + 3x - 5)^3} dx = \int \frac{1}{u^3} \cdot \frac{1}{3} du = \frac{1}{3} \cdot \left(-\frac{1}{2u^2} \right) + C = -\frac{1}{6} \cdot \frac{1}{(x^3 + 3x - 5)^2} + C.$	

Question 5

Skill	Learning Objective	Topic
2.B	FUN-4.A	Determining Concavity of Functions over Their Domains
(A)	Incorrect. These are the intervals for which $f''(x) < 0$, that is, those on which the graph of f is concave down, not concave up. $f'(x) = 12x^2 - 4x^3$ $f''(x) = 24x - 12x^2 = 12x(2 - x)$ The graph of f'' is a parabola opening downward and with zeros at $x = 0$ and $x = 2$. Therefore, $f''(x) < 0$ on the intervals $(-\infty, 0)$ and $(2, \infty)$.	
(B)	Incorrect. The graph of f will be concave up on intervals where $f''(x) > 0$. This response comes from determining where $f'(x) > 0$, however, rather than where $f''(x) > 0$. $f'(x) = 12x^2 - 4x^3 = 4x^2(3 - x) > 0 \text{ when } 3 - x > 0, \text{ so when } x < 3.$	
(C)	Correct. The graph of f will be concave up on intervals where $f''(x) > 0$. $f'(x) = 12x^2 - 4x^3$ $f''(x) = 24x - 12x^2 = 12x(2 - x)$ The graph of f'' is a parabola opening downward and with zeros at $x = 0$ and $x = 2$. Therefore, $f''(x) > 0$ on the interval between the two zeros, or $0 < x < 2$.	
(D)	Incorrect. This response comes from determining where $f(x) > 0$ and $f'(x) > 0$ rather than just where $f''(x) > 0$. $f(x) = 4x^3 - x^4 = x^3(4 - x) > 0 \text{ when } 0 < x < 4.$ $f'(x) = 12x^2 - 4x^3 = 4x^2(3 - x) > 0 \text{ when } x < 3.$ Therefore, $f(x) > 0$ and $f'(x) > 0$ only when $0 < x < 3$.	

Question 6

Skill	Learning Objective	Topic
1.E	FUN-3.D	Implicit Differentiation
(A)	Incorrect. This response would result if there was an error in the power rule when differentiating $3y^{\frac{1}{3}}$ and when solving for $\frac{dy}{dx}$, as follows. $1 + y^{\frac{1}{3}} \frac{dy}{dx} = \frac{dy}{dx}$ At the point $(2, 8)$, $1 + 2 \frac{dy}{dx} = \frac{dy}{dx} \Rightarrow 1 = 3 \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{1}{3}$.	
(B)	Incorrect. This response would result if the chain rule was not used and a sign error was made during the differentiation of the left side, as follows. $1 - y^{-\frac{2}{3}} = \frac{dy}{dx}$ At the point $(2, 8)$, $1 - \frac{1}{4} = \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{3}{4}$.	
(C)	Incorrect. This response would result if the chain rule was not used during the differentiation of the left side, as follows. $1 + y^{\frac{2}{3}} = \frac{dy}{dx}$ At the point $(2, 8)$, $1 + \frac{1}{4} = \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{5}{4}$.	
(D)	Correct. The chain rule is the basis for implicit differentiation. $1 + y^{-\frac{2}{3}} \frac{dy}{dx} = \frac{dy}{dx}$ The point $(2, 8)$ is on the curve since $x = 2$ and $y = 8$ satisfy the equation. At this point, $1 + \frac{1}{4} \frac{dy}{dx} = \frac{dy}{dx} \Rightarrow \frac{3}{4} \frac{dy}{dx} = 1 \Rightarrow \frac{dy}{dx} = \frac{4}{3}$.	

Question 7

Skill	Learning Objective	Topic
1.E	FUN-3.A	Derivatives of $\cos x$, $\sin x$, e^x , and $\ln x$
(A)	Incorrect. This response is an antiderivative of $x^5 - 5^x$, not the derivative.	
(B)	Incorrect. This response would result if the derivative of a^x was taken to be just a^x rather than $(\ln a)a^x$.	
(C)	Incorrect. This response would result if the power rule was applied to the exponential function 5^x , resulting in the response $x \cdot 5^{x-1}$ rather than using the exponential rule $\frac{d}{dx}a^x = (\ln a)a^x$.	
(D)	Correct. The derivative of x^5 is $5x^4$ by the power rule, and the derivative of the exponential function 5^x is $(\ln 5)5^x$. Therefore, $\frac{d}{dx}(x^5 - 5^x) = 5x^4 - (\ln 5)5^x$.	

Question 8

Skill	Learning Objective	Topic
2.B	LIM-2.D	Connecting Limits at Infinity and Horizontal Asymptotes
(A)	Incorrect. This response might come from treating the problem like the limit of a rational function as x goes to infinity when the numerator and denominator are polynomials of the same degree. If only the coefficients of the x^2 term and the e^x term are considered, it might be thought that the limit would be $\frac{-6}{3} = -2$.	
(B)	Correct. The numerator of $\frac{10 - 6x^2}{5 + 3e^x}$ is a translated power function and the denominator is a translated exponential function. Since the exponential function e^x grows faster than the power function x^2 , the relative magnitude of the denominator compared to the numerator will result in this expression converging to 0 as x goes to infinity.	
(C)	Incorrect. This response might come from treating the problem like the limit of a rational function as x goes to 0. If only the constant terms are considered, it might be thought that the limit would be $\frac{10}{5} = 2$.	
(D)	Incorrect. It might be thought that the limit is nonexistent since the numerator goes to $-\infty$ and the denominator goes to $+\infty$ as x goes to infinity, but this does not take into account the relative magnitude of the exponential function in the denominator compared to the quadratic term in the numerator as x gets larger.	

Question 9

Skill	Learning Objective	Topic
1.E	CHA-5.A	Finding the Area Between Curves Expressed as Functions of x
(A)	Incorrect. This response might come from several errors in the antidifferentiation and evaluation. If the term $2x$ is differentiated rather than antidifferentiated, and if in the resulting evaluation the 2 is not included in the evaluation at the endpoints, the result would be as follows. $\int_0^2 (4x - x^2 - 2x) dx = \int_0^2 (2x - x^2) dx = 2 - \frac{1}{3}x^3 \Big _0^2 = 2 - \left(\frac{8}{3} - 0\right) = -\frac{2}{3}$ Then either the negative is ignored or the absolute value is taken, since area must be positive.	
(B)	Correct. The graphs of $y = 2x$ and $y = 4x - x^2$ intersect when $x = 0$ and $x = 2$. The graph of $y = 4x - x^2$ lies above the graph $y = 2x$ on the interval $0 \leq x \leq 2$. (One way to see this is to sketch a graph of the parabola and the line, observing that the graph of $y = 4x - x^2$ has a slope of 4 at $x = 0$, while the graph of $y = 2x$ has a slope of 2.) The area of the region bounded by the two graphs is therefore $\int_0^2 (4x - x^2 - 2x) dx = \int_0^2 (2x - x^2) dx = \left(x^2 - \frac{x^3}{3}\right) \Big _0^2 = 4 - \frac{8}{3} = \frac{4}{3}.$	
(C)	Incorrect. This response might come from two different errors. This first is not finding where the two graphs intersect but looking only at the zeros of $y = 4x - x^2$ at $x = 0$ and $x = 4$, using those as the limits of integration to get $\int_0^4 (4x - x^2 - 2x) dx = \int_0^4 (2x - x^2) dx = \left(x^2 - \frac{x^3}{3}\right) \Big _0^4 = 16 - \frac{64}{3} = -\frac{16}{3},$ and then ignoring the negative (or taking the absolute value, since area must be positive). The response might also come from integrating only $y = 4x - x^2$ over the interval $0 \leq x \leq 2$, as follows. $\int_0^2 (4x - x^2) dx = \left(2x^2 - \frac{x^3}{3}\right) \Big _0^2 = 8 - \frac{8}{3} = \frac{16}{3}$	
(D)	Incorrect. This response might come from adding the two functions rather than taking the difference between them, as follows. $\int_0^2 (4x - x^2 + 2x) dx = \int_0^2 (6x - x^2) dx = \left(3x^2 - \frac{x^3}{3}\right) \Big _0^2 = 12 - \frac{8}{3} = \frac{28}{3}$	

Question 10

Skill	Learning Objective	Topic
1.E	FUN-3.B	The Quotient Rule
(A)	Correct. The derivative of g is found using the quotient rule. $g'(x) = \frac{2xf'(x) - f'(x)(x^2 + 1)}{(f(x))^2}$ The graph of f is used to determine that $f(2) = 3$ and $f'(2) = \frac{7-3}{3-2} = 4$. Then $g'(2) = \frac{4f(2) - f'(2)(5)}{(f(2))^2} = \frac{(4)(3) - (4)(5)}{9} = -\frac{8}{9}$.	
(B)	Incorrect. This response would result if the numerator of the derivative of the quotient was taken to be the product of the derivatives minus the product of the functions, as follows. $g'(x) = \frac{2xf'(x) - f(x)(x^2 + 1)}{(f(x))^2} \Rightarrow g'(2) = \frac{4f'(2) - f(2)(5)}{(f(2))^2} = \frac{(4)(4) - (3)(5)}{9} = \frac{1}{9}$	
(C)	Incorrect. This response would result if the derivative of a quotient was taken to be the quotient of the derivatives, as follows. $g'(x) = \frac{2x}{f'(x)} \Rightarrow g'(2) = \frac{4}{f'(2)} = \frac{4}{4} = 1$	
(D)	Incorrect. This response would result if the terms in the numerator were added rather than subtracted in the quotient rule, as follows. $g'(x) = \frac{2xf'(x) + f'(x)(x^2 + 1)}{(f(x))^2} \Rightarrow g'(2) = \frac{4f(2) + f'(2)(5)}{(f(2))^2} = \frac{(4)(3) + (4)(5)}{9} = \frac{32}{9}$	

Question 11

Skill	Learning Objective	Topic
1.E	FUN-6.A	Applying Properties of Definite Integrals
(A)	Correct. The function $f(x) = \frac{x^2 - x}{x}$ has a removable discontinuity at $x = 0$, since $f(0)$ is undefined but $\lim_{x \rightarrow 0} \frac{x^2 - x}{x} = \lim_{x \rightarrow 0} (x - 1) = -1$. The definition of the definite integral can be extended to functions with removable discontinuities. If $g(x) = x - 1$, then $g(x) = f(x)$ for all x except $x = 0$, and therefore $\int_{-1}^1 f(x) dx = \int_{-1}^1 g(x) dx = \int_{-1}^1 (x - 1) dx = \left(\frac{1}{2}x^2 - x \right) \Big _{-1}^1 = \left(\frac{1}{2} - 1 \right) - \left(\frac{1}{2} + 1 \right) = -2.$	
(B)	Incorrect. This response might arise from an assumption that the value of the definite integral is 0 because the integration is over the symmetric interval $[-1, 1]$.	
(C)	Incorrect. The antiderivative of a quotient might have been taken to be the quotient of antiderivatives, as follows. $\int_{-1}^1 \frac{x^2 - x}{x} dx = \frac{\frac{1}{3}x^3 - \frac{1}{2}x^2}{\frac{1}{2}x^2} \Big _{-1}^1 = \frac{\frac{1}{3} - \frac{1}{2}}{\frac{1}{2}} - \frac{-\frac{1}{3} - \frac{1}{2}}{\frac{1}{2}} = \frac{\frac{2}{3}}{\frac{1}{2}} = \frac{4}{3}$	
(D)	Incorrect. This response might come from not recognizing that the definition of the definite integral can be extended to functions with removable discontinuities.	

Question 12

Skill	Learning Objective	Topic
3.D	FUN-1.B	Using the Mean Value Theorem
(A)	Incorrect. This interval might be chosen because of an error in computing the average rate of change over the interval as $\frac{f(0) - f(4)}{4} = \frac{8 - 0}{4} = 2 \text{ rather than } \frac{f(0) - f(4)}{0 - 4} = \frac{8 - 0}{-4} = -2.$	
(B)	Incorrect. This interval might be chosen because of an error in computing the average rate of change over the interval as $\frac{8 - 4}{f(8) - f(4)} = \frac{4}{2 - 0} = 2 \text{ rather than } \frac{f(8) - f(4)}{8 - 4} = \frac{2 - 0}{4} = \frac{1}{2}.$	
(C)	Correct. The function f is continuous on the closed interval $[8, 12]$ and differentiable on the open interval $(8, 12)$. By the Mean Value Theorem, there is a number c in the interval $(8, 12)$ such that $f'(c) = \frac{f(12) - f(8)}{12 - 8} = \frac{10 - 2}{4} = 2.$	
(D)	Incorrect. This response would result if the Intermediate Value Theorem was used instead of the Mean Value Theorem to select the open interval $(12, 16)$ where $f(c) = 2$ for some number c in the interval.	

Question 13

Skill	Learning Objective	Topic
1.E	LIM-4.A	Using L'Hospital's Rule for Finding Limits of Indeterminate Forms
(A)	Correct. Since $\lim_{x \rightarrow 0} \sin x = 0$ and $\lim_{x \rightarrow 0} (e^x - 1) = 0$, the indeterminate limit can be evaluated using L'Hospital's Rule, as follows. $\lim_{x \rightarrow 0} \frac{\sin x}{e^x - 1} = \lim_{x \rightarrow 0} \frac{\cos x}{e^x} = \frac{\cos 0}{e^0} = \frac{1}{1} = 1$	
(B)	Incorrect. While using L'Hospital's Rule, an error might have been made in the differentiation of e^x, treating it as if taking the derivative of ex, as follows. $\lim_{x \rightarrow 0} \frac{\sin x}{e^x - 1} = \lim_{x \rightarrow 0} \frac{\cos x}{e} = \frac{\cos 0}{e} = \frac{1}{e}$	
(C)	Incorrect. This response would result if the limit of the numerator was observed to be 0, but the denominator was not taken into consideration.	
(D)	Incorrect. This response would result if the limit of the denominator was observed to be 0 while the numerator was not taken into consideration, leading to the assumption that the limit does not exist.	

Question 14

Skill	Learning Objective	Topic
2.C	FUN-7.F	Exponential Models with Differential Equations
(A)	Incorrect. The expression $12t$ might have been treated as if it was the velocity, not the acceleration. Therefore, the position was taken to be $s(t) = 6t^2 + C$, with $C = 5$ since $s(0) = 5$.	
(B)	Incorrect. This response might come from attempting to use the formula $s = \frac{1}{2}at^2 + v_0t + s_0$ for the position of an object falling with constant acceleration a , as follows. $s = \frac{1}{2}at^2 + v_0t + s_0 = \frac{1}{2}(12t)t^2 + 2t + 5 = 6t^3 + 2t + 5$	
(C)	Incorrect. The initial velocity might not have been considered during the antidifferentiation of the acceleration, as follows. $a(t) = 12t \Rightarrow v(t) = 6t^2$ $v(t) = 6t^2 \Rightarrow s(t) = 2t^3 + C;$ $s(0) = 5 \Rightarrow 5 = 0 + 0 + C \Rightarrow C = 5$	
(D)	Correct. Since the acceleration is given, the position can be found using antidifferentiation and the values of the velocity and position at time $t = 0$. $a(t) = 12t \Rightarrow v(t) = 6t^2 + C_1; v(0) = 2 \Rightarrow 2 = 0 + C_1 \Rightarrow C_1 = 2$ $v(t) = 6t^2 + 2 \Rightarrow s(t) = 2t^3 + 2t + C_2;$ $s(0) = 5 \Rightarrow 5 = 0 + 0 + C_2 \Rightarrow C_2 = 5$ The position of the particle is $s(t) = 2t^3 + 2t + 5$ for $t \geq 0$.	

Question 15

Skill	Learning Objective	Topic
3.C	FUN-2.A	Connecting Differentiability and Continuity - Determining When Derivatives Do and Do Not Exist
(A)	Correct. This statement is true. $\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^-} (-x^2 + 3) = -25 + 3 = -22$ $\lim_{x \rightarrow 5^+} f(x) = \lim_{x \rightarrow 5^+} (-10x + 28) = -50 + 28 = -22$ Therefore, $\lim_{x \rightarrow 5} f(x)$ exists and $\lim_{x \rightarrow 5} f(x) = -22 = f(5)$, so f is continuous at $x = 5$. $\lim_{h \rightarrow 0^-} \frac{f(5+h) - f(5)}{h} = \lim_{h \rightarrow 0^-} \frac{-(5+h)^2 + 3 - (-22)}{h} = \lim_{h \rightarrow 0^-} \frac{-10h - h^2}{h} = -10$ $\lim_{h \rightarrow 0^+} \frac{f(5+h) - f(5)}{h} = \lim_{h \rightarrow 0^+} \frac{-10(5+h) + 28 - (-22)}{h} = \lim_{h \rightarrow 0^+} \frac{-10h}{h} = -10$ Therefore, f is also differentiable at $x = 5$ and $f'(5) = -10$. An alternative way to see that the piecewise-defined function f is differentiable at $x = 5$ is to observe that $f'(x) = \begin{cases} -2x & \text{for } x < 5 \\ -10 & \text{for } x > 5. \end{cases}$ Since f is continuous at $x = 5$ and the derivatives $-2x$ and -10 are equal at $x = 5$, f is differentiable at $x = 5$.	
(B)	Incorrect. This statement is false. The function f is both continuous and differentiable at $x = 5$ because $\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^+} f(x) = f(5) = -22$ and $\lim_{h \rightarrow 0^-} \frac{f(5+h) - f(5)}{h} = \lim_{h \rightarrow 0^+} \frac{f(5+h) - f(5)}{h} = -10.$	
(C)	Incorrect. This statement is false, since if f is differentiable at $x = 5$, it must also be continuous at $x = 5$.	
(D)	Incorrect. This statement is false. The function f is both continuous and differentiable at $x = 5$ because $\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^+} f(x) = f(5) = -22$ and $\lim_{h \rightarrow 0^-} \frac{f(5+h) - f(5)}{h} = \lim_{h \rightarrow 0^+} \frac{f(5+h) - f(5)}{h} = -10.$	

Question 16

Skill	Learning Objective	Topic
1.E	FUN-7.D	Finding General Solutions Using Separation of Variables
(A)	Incorrect. This response would result if a chain rule error was made during the antidifferentiation of the dy term. $\frac{dy}{dx} = 2 - y \Rightarrow \frac{dy}{2 - y} = dx$ $\int \frac{1}{2 - y} dy = \int dx \Rightarrow \ln 2 - y = x + C$ $\ln 1 = 1 + C \Rightarrow C = -1$ $\ln 2 - y = x - 1 \Rightarrow 2 - y = e^{x-1}$ Since $2 - y > 0$ at the initial value $y = 1$, the solution would be $2 - y = e^{x-1}$, or $y = 2 - e^{x-1}$.	
(B)	Correct. The differential equation can be solved using separation of variables and the initial condition to determine the appropriate value for the arbitrary constant. $\frac{dy}{dx} = 2 - y \Rightarrow \frac{dy}{2 - y} = dx$ $\int \frac{1}{2 - y} dy = \int dx \Rightarrow -\ln 2 - y = x + C$ $-\ln 1 = 1 + C \Rightarrow C = -1$ $-\ln 2 - y = x - 1 \Rightarrow \ln 2 - y = -x + 1 \Rightarrow 2 - y = e^{1-x}$ Since $2 - y > 0$ at the initial value $y = 1$, the solution to the differential equation is $2 - y = e^{1-x}$, or $y = 2 - e^{1-x}$.	
(C)	Incorrect. This response would result if an arbitrary constant was not included during the antidifferentiation. $\frac{dy}{dx} = 2 - y \Rightarrow \frac{dy}{2 - y} = dx$ $\int \frac{1}{2 - y} dy = \int dx \Rightarrow -\ln 2 - y = x \Rightarrow 2 - y = e^{-x}$ Since $2 - y > 0$ at the initial value $y = 1$, the solution would be $2 - y = e^{-x}$, or $y = 2 - e^{-x}$.	
(D)	Incorrect. This response would result if an arbitrary constant was not included during the antidifferentiation and the incorrect sign was taken for the absolute value when solving for y. $\frac{dy}{dx} = 2 - y \Rightarrow \frac{dy}{2 - y} = dx$ $\int \frac{1}{2 - y} dy = \int dx \Rightarrow -\ln 2 - y = x \Rightarrow 2 - y = e^{-x}$ $2 - y = -e^{-x} \Rightarrow y = 2 + e^{-x}$	

Question 17

Skill	Learning Objective	Topic
2.B	FUN-5.A	Interpreting the Behavior of Accumulation Functions Involving Area
(A)	Incorrect. This is the x -coordinate of a critical point of g , not of a point of inflection of the graph of g . The equation $g'(x) = x^2 - 5x - 14 = (x + 2)(x - 7) = 0$ might have been solved for x rather than the equation $g''(x) = 2x - 5 = 0$.	
(B)	Correct. To find the point of inflection of the graph of g, determine where g'' changes sign. $g'(x) = x^2 - 5x - 14$ $g''(x) = 2x - 5$ Then $g''(x) = 0$ at $x = \frac{5}{2}$. Since $g''(x) < 0$ for $x < \frac{5}{2}$ and $g''(x) > 0$ for $x > \frac{5}{2}$, the graph of g changes concavity at $x = \frac{5}{2}$ and therefore, the graph of g has a point of inflection at $x = \frac{5}{2}$.	
(C)	Incorrect. This is a value of x where $g(x) = 0$, not a value where $g''(x) = 0$.	
(D)	Incorrect. This is the x -coordinate of a critical point of g , not of a point of inflection of the graph of g . The equation $g'(x) = x^2 - 5x - 14 = (x + 2)(x - 7) = 0$ might have been solved for x rather than the equation $g''(x) = 2x - 5 = 0$.	

Question 18

Skill	Learning Objective	Topic
1.E	FUN-3.B	The Product Rule
(A)	Incorrect. This response would result if the chain rule was correctly used for the derivative of $\sec(2x)$ but the product rule was incorrectly applied as $\frac{d}{dx}(f(x)g(x)) = f'(x)g'(x)$. $3x^2 \cdot (2\sec(2x)\tan(2x))$	
(B)	Incorrect. This response would result if the product rule was correctly used but the derivative of $\sec(2x)$ was taken to be $2\tan^2(2x)$. $x^3 \cdot \frac{d}{dx}(\sec(2x)) + 3x^2 \cdot \sec(2x) = x^3 \cdot (\tan^2(2x) \cdot 2) + 3x^2\sec(2x)$	
(C)	Incorrect. This response would result if the product rule was correctly used but the chain rule was not used for the derivative of $\sec(2x)$. $x^3 \cdot \frac{d}{dx}(\sec(2x)) + 3x^2 \cdot \sec(2x) = x^3 \cdot (\sec(2x)\tan(2x)) + 3x^2\sec(2x)$	
(D)	Correct. A combination of the product rule and the chain rule is used to compute the derivative. $\frac{d}{dx}(x^3\sec(2x)) = x^3 \cdot \frac{d}{dx}(\sec(2x)) + 3x^2 \cdot \sec(2x)$ $= x^3 \cdot (\sec(2x)\tan(2x) \cdot 2) + 3x^2\sec(2x)$ $= 2x^3\sec(2x)\tan(2x) + 3x^2\sec(2x)$	

Question 19

Skill	Learning Objective	Topic
2.B	CHA-3.B	Straight-Line Motion: Connecting Position, Velocity, and Acceleration
(A)	Correct. The velocity of the particle is $v(t) = 3t^2 - 8t + 4$. At time $t = 1$, $v(1) = -1$. Since the velocity is negative, the particle is moving down the y -axis. The rate of change of the velocity is $v'(t) = 6t - 8$. At time $t = 1$, $v'(1) = -2$. Since this is negative, the particle is moving with decreasing velocity at time $t = 1$.	
(B)	Incorrect. It was correctly determined that the particle is moving down the y -axis, but since $v'(t) = 6t - 8$ and $v'(1) = -2$, the particle's velocity is decreasing, not increasing.	
(C)	Incorrect. It was correctly determined that the particle is moving with decreasing velocity, but since $v(t) = 3t^2 - 8t + 4$ and $v(1) = -1$, the particle is moving down the y -axis, not up the axis.	
(D)	Incorrect. Since $v(t) = 3t^2 - 8t + 4$ and $v(1) = -1$, the particle is moving down the y -axis, not up the axis. Since $v'(t) = 6t - 8$ and $v'(1) = -2$, the particle's velocity is decreasing, not increasing.	

Question 20

Skill	Learning Objective	Topic
1.C	FUN-6.A	Applying Properties of Definite Integrals
(A)	Incorrect. The value of this integral can be determined using the properties of the definite integral, as follows. $\int_4^1 g(x) \, dx = -\int_1^4 g(x) \, dx = -(-2) = 2$	
(B)	Incorrect. The value of this integral can be determined using the properties of the definite integral, as follows. $\int_1^4 3f(x) \, dx = 3 \cdot \int_1^4 f(x) \, dx = 3 \cdot 8 = 24$	
(C)	Correct. It is not true in general that $\int_1^4 3f(x)g(x) \, dx = \int_1^4 3f(x) \, dx \cdot \int_1^4 g(x) \, dx$, so the individual values of $\int_1^4 f(x) \, dx$ and $\int_1^4 g(x) \, dx$ cannot be used to determine the value of $\int_1^4 3f(x)g(x) \, dx$. For example, if $f(x) = \frac{8}{3}$ and $g(x) = -\frac{2}{3}$, then $\int_1^4 f(x) \, dx = 8$, $\int_1^4 g(x) \, dx = -2$, and $\int_1^4 3f(x)g(x) \, dx = \int_1^4 -\frac{16}{3} \, dx = -16$. However, if $f(x) = \frac{16}{9}(x-1)$ and $g(x) = -\frac{4}{9}(x-1)$, then $\int_1^4 f(x) \, dx = 8$ and $\int_1^4 g(x) \, dx = -2$ as before, but now $\int_1^4 3f(x)g(x) \, dx = \int_1^4 -\frac{64}{27}(x-1)^2 \, dx = -\frac{64}{3}$.	
(D)	Incorrect. The value of this integral can be determined using the properties of the definite integral, as follows. $\begin{aligned} \int_1^4 (3f(x) + g(x)) \, dx &= \int_1^4 3f(x) \, dx + \int_1^4 g(x) \, dx \\ &= 3 \cdot \int_1^4 f(x) \, dx + \int_1^4 g(x) \, dx = 3 \cdot 8 + (-2) = 22 \end{aligned}$	

Question 21

Skill	Learning Objective	Topic
1.D	CHA-2.B	Defining the Derivative of a Function and Using Derivative Notation
(A)	Incorrect. This response might come from observing that the numerator is zero when $h = 0$ without consideration of the denominator.	
(B)	Incorrect. This response might come from observing that both the numerator and the denominator are zero when $h = 0$ and interpreting $\frac{0}{0}$ as equal to 1.	
(C)	Incorrect. This response would result if the limit of the difference quotient was correctly recognized as the derivative of the function $f(x) = \sin(2x)$, but the chain rule was not used in finding the derivative.	
(D)	Correct. The limit of this difference quotient is of the form $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$, where $f(x) = \sin(2x)$. This is one way to express the derivative of f . By the chain rule, $f'(x) = \cos(2x) \cdot 2$.	

Question 22

Skill	Learning Objective	Topic
2.D	FUN-7.C	Sketching Slope Fields
(A)	Incorrect. This response might be chosen because the slopes for the differential equation $\frac{dy}{dx} = x + y $ are nonnegative, which matches the behavior shown in the slope field. However, the segments in the slope field where $y = 0$ all have slope 0 and that would not be true for this differential equation.	
(B)	Incorrect. This response might be chosen because the slope segments look like they might match the behavior of the cubic polynomial $f(x) = x^3$. However, the slopes of the segments in the slope field depend only on the variable y and that would not be true for the differential equation $\frac{dy}{dx} = x^3$.	
(C)	Incorrect. This response might be chosen because the slopes for the differential equation $\frac{dy}{dx} = y^3$ depend only on the variable y and have the value 0 when $y = 0$, which matches the behavior shown in the slope field. However, the segments in the slope field all have nonnegative slopes and that would not be true for this differential equation when $y < 0$.	
(D)	Correct. The segments in the slope field suggest that (1) the slopes depend only on the variable y , (2) the slopes are nonnegative, and (3) the slopes are zero when $y = 0$. The differential equation $\frac{dy}{dx} = y^2$ satisfies all three conditions. The differential equation $\frac{dy}{dx} = x + y $ does not satisfy conditions (1) and (3). The differential equation $\frac{dy}{dx} = x^3$ does not satisfy any of the conditions. The differential equation $\frac{dy}{dx} = y^3$ does not satisfy condition (2) when $y < 0$.	

Question 23

Skill	Learning Objective	Topic
1.E	CHA-5.B	Volumes with Cross Sections - Squares and Rectangles
(A)	Incorrect. This response is the area of the region, not the volume of the solid. $\int_0^1 e^x dx = e^x \Big _0^1 = e - 1$	
(B)	Correct. The area of a square of side length s is s^2. A typical cross section of the solid is a square with side from the x-axis to the graph of $y = e^x$. The length of the side of the square is therefore $s = e^x$, so the area of the square is $(e^x)^2 = e^{2x}$. The volume of the solid is found using the definite integral of the cross-sectional area. $\int_0^1 e^{2x} dx = \frac{1}{2} e^{2x} \Big _0^1 = \frac{1}{2} e^2 - \frac{1}{2}$	
(C)	Incorrect. This response would result if the volume was set up correctly as the definite integral of the cross-sectional area e^{2x}, but an error was made in the antidifferentiation by not considering the chain rule, as follows. $\int_0^1 e^{2x} dx = e^{2x} \Big _0^1 = e^2 - 1$	
(D)	Incorrect. This response would result if the volume was set up correctly as the definite integral of the cross-sectional area e^{2x}, but an error was made with respect to the chain rule in the antidifferentiation of the exponential function (or the integrand was differentiated rather than antidifferentiated), as follows. $\int_0^1 e^{2x} dx = 2e^{2x} \Big _0^1 = 2e^2 - 2$	

Question 24

Skill	Learning Objective	Topic
3.D	FUN-6.B	The Fundamental Theorem of Calculus and Definite Integrals
(A)	Correct. By the Fundamental Theorem of Calculus, $\int_0^{12} f'(x) \, dx = f(12) - f(0) = (-4) - 4 = -8,$ where the values of f at $x = 0$ and $x = 12$ are obtained from the graph.	
(B)	Incorrect. This response would result if the function f was integrated over the interval $[0, 12]$ rather than f', as follows. $\int_0^{12} f(x) \, dx = \frac{1}{2}(4)(4) - \frac{1}{2}(4)(3) - \frac{1}{2}(4)(4) = -6$	
(C)	Incorrect. This response would result if the Fundamental Theorem of Calculus was incorrectly applied, as follows. $\int_0^{12} f'(x) \, dx = f'(12) - f'(0) = (-1) - (-1) = 0$	
(D)	Incorrect. This response is the total area bounded by the graph of f and the x-axis over the interval $[0, 12]$. $\int_0^{12} f(x) \, dx = \frac{1}{2}(4)(4) + \frac{1}{2}(4)(3) + \frac{1}{2}(4)(4) = 8 + 6 + 8 = 22$	

Question 25

Skill	Learning Objective	Topic
3.G	FUN-7.B	Verifying Solutions for Differential Equations
(A)	Correct. One way to verify that a function is a solution to a differential equation is to check that the function and its derivatives satisfy the differential equation. The differential equation in this option involves y and y'. The correct derivative must be computed and the algebra correctly done to verify that the differential equation is satisfied. $y' = 12e^{6x}$ $y' - 6y - 30 = 12e^{6x} - 6(2e^{6x} - 5) - 30 = 12e^{6x} - 12e^{6x} + 30 - 30 = 0$	
(B)	Incorrect. The correct derivative was found, but this differential equation will appear to be satisfied if the 12 in the second term is not distributed correctly across the two terms in the parentheses, as follows. $y' = 12e^{6x}$ $2y' - 12y + 5 = 24e^{6x} - 12(2e^{6x} - 5) + 5 = 24e^{6x} - 24e^{6x} + 5 - 5 = 0$	
(C)	Incorrect. This differential equation has $y = 2e^{6x}$ as a solution, not $y = 2e^{6x} - 5$. Both functions have $y' = 12e^{6x}$ and $y'' = 72e^{6x}$, but $72e^{6x} - 5(12e^{6x}) - 6(2e^{6x}) = 0$, whereas $72e^{6x} - 5(12e^{6x}) - 6(2e^{6x} - 5) = 30$.	
(D)	Incorrect. This differential equation will appear to be satisfied if the chain rule is not used in taking the derivative of the exponential, as follows. $y' = 2e^{6x} \quad y'' = 2e^{6x}$ $y'' - 2y' + y + 5 = 2e^{6x} - 2(2e^{6x}) + (2e^{6x} - 5) + 5 = 2e^{6x} - 4e^{6x} + 2e^{6x} - 5 + 5 = 0$	

Question 26

Skill	Learning Objective	Topic
3.D	FUN-4.A	Using the Candidates Test to Find Absolute (Global) Extrema
(A)	Incorrect. This response would result if the critical point was not found, and the endpoint with the smallest function value was selected.	
(B)	Correct. The absolute minimum will occur at a critical point or one of the endpoints. $y' = 4x^{\frac{1}{3}} - 2 = 0 \Rightarrow x^{\frac{1}{3}} = \frac{1}{2} \Rightarrow x = \left(\frac{1}{2}\right)^3 = \frac{1}{8}$ The candidates are $x = 0$, $x = \frac{1}{8}$, and $x = 1$. When $x = 0$, $y = 0$. When $x = \frac{1}{8}$, $y = 3\left(\frac{1}{8}\right)^{\frac{4}{3}} - 2\left(\frac{1}{8}\right) = \frac{3}{16} - \frac{1}{4} = -\frac{1}{16}$. When $x = 1$, $y = 1$. The absolute minimum is therefore at $x = \frac{1}{8}$. Alternatively, since $x = \frac{1}{8}$ is the only critical point and the Second Derivative Test shows that it is the location of a local minimum, it must also be the location of the absolute minimum on the interval $[0, 1]$.	
(C)	Incorrect. This response would result if an error in the power rule has the derivative being computed as $y' = 4x - 2$ with a zero at $x = \frac{1}{2}$. It was concluded that this was the location of a local minimum from the Second Derivative Test and therefore had to be the location of the absolute minimum, since it was the only critical point in the interval $[0, 1]$. (Note that $y\left(\frac{1}{2}\right) = 3\left(\frac{1}{16}\right)^{\frac{1}{3}} - 1$ and it is not obvious whether this value is smaller or greater than $y(0) = 0$, so it would have been difficult to use the Candidates Test. The y-value at $x = \frac{1}{2}$ is actually positive.)	
(D)	Incorrect. This is the value of x at which the maximum value of y occurs on the interval $[0, 1]$.	

Question 27

Skill	Learning Objective	Topic
3.F	CHA-3.A	Interpreting the Meaning of the Derivative in Context
(A)	Incorrect. The rate at which the depth of the water is increasing is $W'(t)$. Therefore, this sentence is an interpretation of the statement $W(2) > 3$, not an interpretation about $W'(t)$.	
(B)	Incorrect. The rate at which the depth of the water is increasing is $W'(t)$. Therefore, this sentence is an interpretation of the statement $W'(t) > 3$ for all t in the interval $0 \leq t \leq 2$.	
(C)	Correct. In the expression $W'(2)$, the 2 represents the value of the independent variable and is therefore the number of hours since the tank began filling with water. $W'(2)$, being the value of a derivative, is the rate of change of W , that is, the rate of change of the rate at which the depth of the water is rising; in this case, 2 hours after the tank begins filling with water. The units for the derivative would be the units of W per unit of time; thus, feet per hour per hour. The statement says that at time 2 hours after the tank begins filling with water, the rate at which the depth of the water is rising, $W'(t)$, is increasing at a rate that is greater than 3 feet per hour per hour.	
(D)	Incorrect. This sentence is an interpretation of the statement $W'(t) > 3$ for all t in the interval $0 \leq t \leq 2$. $W'(2)$, being the value of a derivative, is the instantaneous rate of change of W at the particular instant $t = 2$.	

Question 28

Skill	Learning Objective	Topic
1.E	CHA-3.E	Solving Related Rates Problems
(A)	Incorrect. This response would result if the relationship $\tan \theta = \frac{h}{30}$ was correctly used, where h is the height of the balloon above the point P at time t, but then the derivative of $\tan \theta$ was taken to be $\sec \theta \tan \theta$ rather than $\sec^2 \theta$. $\sec \theta \tan \theta \frac{d\theta}{dt} = \frac{1}{30} \frac{dh}{dt}$ At the instant when $h = 40$, the distance from the person to the balloon is $\sqrt{30^2 + 40^2} = \sqrt{2500} = 50$. $\left(\frac{50}{30}\right)\left(\frac{40}{30}\right) \frac{d\theta}{dt} = \frac{1}{30} \cdot 2 = \frac{1}{15} \Rightarrow \frac{d\theta}{dt} = \frac{1}{15} \cdot \frac{3}{5} \cdot \frac{3}{4} = \frac{3}{100}$	
(B)	Correct. If $h(t)$ is the height of the balloon above the point P at time t, then $\tan \theta = \frac{h}{30}$. Using implicit differentiation with respect to t shows that $\sec^2 \theta \frac{d\theta}{dt} = \frac{1}{30} \frac{dh}{dt}$. At the instant when $h = 40$, the distance from the person to the balloon is $\sqrt{30^2 + 40^2} = \sqrt{2500} = 50. \text{ At this instant, } \sec \theta = \frac{50}{30} = \frac{5}{3} \text{ and}$ therefore $\left(\frac{5}{3}\right)^2 \frac{d\theta}{dt} = \frac{1}{30} \cdot 2 = \frac{1}{15} \Rightarrow \frac{d\theta}{dt} = \frac{1}{15} \cdot \frac{9}{25} = \frac{3}{125}$.	
(C)	Incorrect. This response would result if the relationship $\tan \theta = \frac{h}{30}$ was correctly used, where h is the height of the balloon above the point P at time t, but then the derivative of $\tan \theta$ was taken to be $\sin \theta$ rather than $\sec^2 \theta$. $\sin \theta \frac{d\theta}{dt} = \frac{1}{30} \frac{dh}{dt}$ At the instant when $h = 40$, the distance from the person to the balloon is $\sqrt{30^2 + 40^2} = \sqrt{2500} = 50$. $\left(\frac{40}{50}\right) \frac{d\theta}{dt} = \frac{1}{30} \cdot 2 = \frac{1}{15} \Rightarrow \frac{d\theta}{dt} = \frac{1}{15} \cdot \frac{5}{4} = \frac{1}{12}$	
(D)	Incorrect. This response would result if the relationship $\tan \theta = \frac{h}{30}$ was correctly used, where h is the height of the balloon above the point P at time t, but then the derivative of $\tan \theta$ was taken to be $\cos^2 \theta$ rather than $\sec^2 \theta$. $\cos^2 \theta \frac{d\theta}{dt} = \frac{1}{30} \frac{dh}{dt}$ At the instant when $h = 40$, the distance from the person to the balloon is $\sqrt{30^2 + 40^2} = \sqrt{2500} = 50$. $\left(\frac{30}{50}\right)^2 \frac{d\theta}{dt} = \frac{1}{30} \cdot 2 = \frac{1}{15} \Rightarrow \frac{d\theta}{dt} = \frac{1}{15} \cdot \frac{25}{9} = \frac{5}{27}$	

Question 29

Skill	Learning Objective	Topic
2.B	LIM-2.D	Connecting Infinite Limits and Vertical Asymptotes
(A)	Correct. Since $\frac{x-2}{x^4-16} = \frac{x-2}{(x^2-4)(x^2+4)} = \frac{x-2}{(x-2)(x+2)(x^2+4)} = \frac{1}{(x+2)(x^2+4)}$ for $x \neq 2$, the graph has only one vertical asymptote, which is at $x = -2$. The graph has a removable discontinuity at $x = 2$.	
(B)	Incorrect. The denominator is $x^4 - 16 = (x^2 - 4)(x^2 + 4) = (x - 2)(x + 2)(x^2 + 4)$ which has two zeros, which might lead to the conclusion that there are two vertical asymptotes.	
(C)	Incorrect. The expression might have been correctly simplified to $\frac{x-2}{x^4-16} = \frac{x-2}{(x^2-4)(x^2+4)} = \frac{x-2}{(x-2)(x+2)(x^2+4)} = \frac{1}{(x+2)(x^2+4)}$ for $x \neq 2$, but then the conclusion was made that there are three vertical asymptotes because the denominator is a cubic polynomial.	
(D)	Incorrect. Since the denominator is a quartic polynomial, the assumption might have been made that there are four zeros and therefore four vertical asymptotes.	

Question 30

Skill	Learning Objective	Topic
3.D	LIM-5.C	Riemann Sums, Summation Notation, and Definite Integral Notation
(A)	Incorrect. The sum can be interpreted as a right Riemann sum in the form $\sum_{k=1}^n f(1 + k\Delta x)\Delta x$, where $f(x) = x^2$ and $\Delta x = \frac{2}{n}$. The value of Δx corresponds to an interval of length 2, but b is not equal to 2 because the interval starts at $x = 1$.	
(B)	Correct. The sum can be interpreted as a right Riemann sum in the form $\sum_{k=1}^n f(1 + k\Delta x)\Delta x$, where $f(x) = x^2$ and $\Delta x = \frac{2}{n}$. The value of Δx corresponds to an interval of length 2. The sum starts with the right endpoint $1 + \Delta x$ and ends with the right endpoint $1 + n\Delta x = 1 + 2 = 3$, so the Riemann sum is over the interval $[1, 3]$. The limit of the Riemann sum is the definite integral $\int_1^3 f(x) dx$. There could not be another value of b for which $\int_1^b x^2 dx$ has the same value as $\int_1^3 x^2 dx$ since $I(b) = \int_1^b x^2 dx$ is a strictly increasing function of b . Therefore, $b = 3$ is the only choice.	
(C)	Incorrect. Suppose the value of the limit is A . The equation $A = \int_1^b x^2 dx = \frac{b^3}{3} - \frac{1}{3}$ has only one solution, $b = (3A + 1)^{\frac{1}{3}}$. Therefore, b could not be any real number.	
(D)	Incorrect. This response might come from believing that the limit does not exist since it involves an infinite summation.	

Question 76

Skill	Learning Objective	Topic
2.E	FUN-4.A	Determining Intervals on Which a Function Is Increasing or Decreasing
(A)	Incorrect. The graph of f is concave up where f' is increasing. This response might come from switching the roles of the function and its derivative and thinking that f is increasing where the graph of f' is concave up. The graph of f' is concave up on the intervals $(0, 1)$ and $(2, 4)$.	
(B)	Incorrect. This response might come from treating the given graph as the graph of f rather than the graph of f' . These are the two intervals where f' is increasing.	
(C)	Incorrect. These are the intervals where both f and f' are increasing.	
(D)	Correct. The function f is increasing on closed intervals where f' is positive on the corresponding open intervals. The graph indicates that $f'(x) > 0$ on the intervals $(0, 2)$ and $(4, 5)$, so f is increasing on the intervals $[0, 2]$ and $[4, 5]$.	

Question 77

Skill	Learning Objective	Topic
1.E	CHA-3.B	Straight-Line Motion: Connecting Position, Velocity, and Acceleration
(A)	Incorrect. This response is the acceleration of the object at time $t = 3$.	
(B)	Correct. The velocity is the derivative of the height. Using the calculator, $v(3) = h'(3) = 7.778$.	
(C)	Incorrect. This response is the height of the object at time $t = 3$.	
(D)	Incorrect. This response is the value of $\int_1^3 h(t) dt$.	

Question 78

Skill	Learning Objective	Topic
1.E	CHA-3.F	Approximating Values of a Function Using Local Linearity and Linearization
(A)	Correct. An equation of the line tangent to the graph of g at $x = a$ is $y = g(a) + g'(a)(x - a)$. In this question, $a = -1$. The value of y when $x = -1.2$ would be an approximation to $g(-1.2)$. $g(-1.2) \approx g(-1) + g'(-1)(-1.2 - (-1)) = 4 + 2(-0.2) = 3.6$	
(B)	Incorrect. This response would result if the derivative was not used as the slope of the tangent line, as follows. $g(-1.2) \approx g(-1) + \Delta x = 4 + (-0.2) = 3.8$	
(C)	Incorrect. This response would result if the derivative was not used as the slope of the tangent line, and Δx was taken to be 0.2 rather than -0.2 , as follows. $g(-1.2) \approx g(-1) + \Delta x = 4 + 0.2 = 4.2$	
(D)	Incorrect. During the evaluation of the change in y along the tangent line, the change in x was incorrectly taken to be 0.2 rather than -0.2 , as follows. $g(-1.2) \approx g(-1) + \Delta y = g(-1) + g'(-1)\Delta x = 4 + 2(0.2) = 4.4$	

Question 79

Skill	Learning Objective	Topic
3.F	CHA-4.B	Finding the Average Value of a Function on an Interval
(A)	Incorrect. The definite integral was not divided by the length of the interval $[30, 60]$ over which the averaging is done.	
(B)	Correct. The average value of a function f over an interval $[a, b]$ is $\frac{1}{b-a} \int_a^b f(x) dx$. Tara's average heart rate from $t = 30$ to $t = 60$ is the average value of the function h over the interval $[30, 60]$ and would therefore be given by the expression $\frac{1}{60-30} \int_{30}^{60} h(t) dt$.	
(C)	Incorrect. This response is the average rate of change of Tara's heart rate from $t = 30$ to $t = 60$, not the average of her heart rate over that interval. By the Fundamental Theorem of Calculus, this expression is equal to $\frac{h(60) - h(30)}{60 - 30}$.	
(D)	Incorrect. This response is the average of the rate of change of Tara's heart rate at the two times $t = 30$ and $t = 60$, not the average of her heart rate over the interval from $t = 30$ to $t = 60$.	

Question 80

Skill	Learning Objective	Topic
1.E	FUN-6.B	The Fundamental Theorem of Calculus and Definite Integrals
(A)	Incorrect. This response comes from taking $g(5) = g(2) + g'(5) = -7 + g'(5) = 4.402$.	
(B)	Incorrect. This response is the value of $g'(5)$, not the value of $g(5)$.	
(C)	Correct. By the Fundamental Theorem of Calculus, $g(5) - g(2) = \int_2^5 g'(x) dx$. Therefore, $g(5) = g(2) + \int_2^5 \sqrt{x^3 + x} dx = -7 + \int_2^5 \sqrt{x^3 + x} dx = 13.899$, where the evaluation of the definite integral is done with the calculator.	
(D)	Incorrect. This response would result if the initial condition was not included in the computation, resulting in $g(5) = \int_2^5 \sqrt{x^3 + x} dx = 20.899$.	

Question 81

Skill	Learning Objective	Topic
2.B	LIM-2.A	Exploring Types of Discontinuities
(A)	Incorrect. The function corresponding to this graph has a jump discontinuity at $x = 3$, not a removable one, because $\lim_{x \rightarrow 3^-} f(x) \neq \lim_{x \rightarrow 3^+} f(x)$.	
(B)	Incorrect. The function corresponding to this graph has a jump discontinuity at $x = 3$, not a removable one, because $\lim_{x \rightarrow 3^-} f(x) \neq \lim_{x \rightarrow 3^+} f(x)$.	
(C)	Correct. A removable discontinuity occurs at $x = c$ if $\lim_{x \rightarrow c} f(x)$ exists, but $f(c)$ does not exist or is not equal to the value of the limit. This graph could be the graph of f since $\lim_{x \rightarrow 3} f(x)$ exists but is not equal to $f(3)$.	
(D)	Incorrect. The function corresponding to this graph has a discontinuity at $x = 3$ due to a vertical asymptote, not a removable discontinuity.	

Question 82

Skill	Learning Objective	Topic
1.E	FUN-6.A	Applying Properties of Definite Integrals
(A)	Correct. Using the property of definite integrals over adjacent intervals, $\int_0^{20} f(x) dx = \int_0^{17} f(x) dx + \int_{17}^{20} f(x) dx = 8 + (-3) = 5.$ Another application of the same property gives $\int_0^{20} f(x) dx = \int_0^{13} f(x) dx + \int_{13}^{20} f(x) dx \Rightarrow \int_0^{13} f(x) dx = \int_0^{20} f(x) dx - \int_{13}^{20} f(x) dx.$ Therefore, $\int_0^{13} f(x) dx = \int_0^{20} f(x) dx - \int_{13}^{20} f(x) dx = 5 - 7 = -2.$	
(B)	Incorrect. Using the property of definite integrals over adjacent intervals, $\int_0^{20} f(x) dx = \int_0^{13} f(x) dx + \int_{13}^{20} f(x) dx \Rightarrow \int_0^{13} f(x) dx = \int_0^{20} f(x) dx - \int_{13}^{20} f(x) dx.$ However, if $\int_0^{20} f(x) dx$ was incorrectly determined to be $\int_0^{17} f(x) dx - \int_{17}^{20} f(x) dx = 8 - (-3) = 11 \text{ rather than}$ $\int_0^{17} f(x) dx + \int_{17}^{20} f(x) dx = 8 + (-3) = 5, \text{ the result would be as follows.}$ $\int_0^{13} f(x) dx = \int_0^{20} f(x) dx - \int_{13}^{20} f(x) dx = 11 - 7 = 4$	
(C)	Incorrect. This response would result if the property of definite integrals over adjacent intervals was not used appropriately. The values of the three definite integrals might have been added, as follows. $\int_0^{17} f(x) dx + \int_{17}^{20} f(x) dx + \int_{13}^{20} f(x) dx = 8 + (-3) + 7 = 12$	
(D)	Incorrect. The property of definite integrals over adjacent intervals was not used appropriately. The absolute values of the definite integrals were added, as follows. $\left \int_0^{17} f(x) dx \right + \left \int_{17}^{20} f(x) dx \right + \left \int_{13}^{20} f(x) dx \right = 8 + 3 + 7 = 18$	

Question 83

Skill	Learning Objective	Topic
3.D	LIM-2.C	Removing Discontinuities
(A)	Correct. The limit at $x = 3$ exists if the left-hand and right-hand limits are equal. $\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x) \Rightarrow k^3 + 3 = \frac{16}{k^2 - 3}$ The solution to this equation for $k > 0$ is $k = 2.081$. With this value of k, $\lim_{x \rightarrow 3} f(x)$ exists and is equal to $f(3)$. Therefore, f is continuous at $x = 3$.	
(B)	Incorrect. This response comes from trying to make the left-hand and right-hand limits of the derivative equal at $x = 3$, as follows. $f'(x) = \begin{cases} 1 & \text{for } x < 3 \\ \frac{16}{(k^2 - x)^2} & \text{for } x > 3 \end{cases}$ $\lim_{x \rightarrow 3^-} f'(x) = \lim_{x \rightarrow 3^+} f'(x) \Rightarrow 1 = \frac{16}{(k^2 - 3)^2}$ The solution to this equation for $k > 0$ is $k = 2.646$.	
(C)	Incorrect. In trying to set the left-hand and right-hand limits of f equal at $x = 3$, the 3 might have been substituted for the parameter k rather than the variable x, as follows. $27 + x = \frac{16}{9 - x}$ The positive solution to this equation is $x = 8.550$.	
(D)	Incorrect. This response might come from errors that lead to an equation that has no positive solution. For example, it might come from trying to make the left-hand and right-hand limits of the derivative equal at $x = 3$ but also making a chain rule error in the derivative of the piece for $x > 3$, as follows. $f'(x) = \begin{cases} 1 & \text{for } x < 3 \\ \frac{-16}{(k^2 - x)^2} & \text{for } x > 3 \end{cases}$ $\lim_{x \rightarrow 3^-} f'(x) = \lim_{x \rightarrow 3^+} f'(x) \Rightarrow 1 = \frac{-16}{(k^2 - 3)^2}$ This equation has no solution for k.	

Question 84

Skill	Learning Objective	Topic
3.D	FUN-1.A	Working with the Intermediate Value Theorem
(A)	Incorrect. By the Intermediate Value Theorem, the function f is guaranteed to have at least one zero, but it could have more. For example, the function $f(x) = \frac{2}{3}x^3 + 2x^2 - \frac{14}{3}x + 1$ satisfies the values in the table, but it has three zeros.	
(B)	Incorrect. The largest value of f in the table occurs at $x = 2$, but that does not guarantee that f has a relative maximum at $x = 2$. For example, the function $f(x) = \frac{2}{3}x^3 + 2x^2 - \frac{14}{3}x + 1$ satisfies the values in the table, but it does not have a relative maximum at $x = 2$ because $f'(2) > 0$.	
(C)	Correct. Since f is continuous on the closed interval $[-5, 2]$ and $f(-5) < 4 < f(2)$, then by the Intermediate Value Theorem there must be a value c in the open interval $(-5, 2)$ such that $f(c) = 4$.	
(D)	Incorrect. While it is true that $\frac{f(2) - f(-5)}{2 - (-5)} = \frac{14}{7} = 2$, the Mean Value Theorem cannot be used to claim that there exists a value c in the open interval $(-5, 2)$ such that $f'(c) = 2$ because there is no assumption that f is differentiable on the open interval $(-5, 2)$. Consider the function f defined as follows. $f(x) = \begin{cases} -9 & \text{for } x \leq -1 \\ 10x + 1 & \text{for } -1 < x \leq 0 \\ 4x + 1 & \text{for } 0 < x \leq 1 \\ 5 & \text{for } x > 1 \end{cases}$ The graph of f goes through each of the points in the table, but none of the linear pieces of the graph has slope 2.	

Question 85

Skill	Learning Objective	Topic
3.D	FUN-4.A	Using the First Derivative Test to Find Relative (Local) Extrema
(A)	Incorrect. This is a value of x where $f'(x) = g(x) = 0$. But since the graph of $y = g(x)$ goes from negative to positive at this point, this would be a local minimum for the graph of $y = f(x)$, not a local maximum.	
(B)	Incorrect. This is a value of x where $g'(x) = 0$, not where $f'(x) = g(x) = 0$. It is the x -coordinate of a local maximum for the graph of $y = g(x)$, not for the graph of $y = f(x)$.	
(C)	Incorrect. This is a value of x where $g'(x) = 0$, not where $f'(x) = g(x) = 0$. It is the x -coordinate of a local minimum for the graph of $y = g(x)$.	
(D)	Correct. A local maximum for the graph of $y = f(x)$ occurs at a value of x where $f' = g$ changes from positive to negative. The graph of $y = g(x)$ crosses the x -axis from positive to negative at $x = 3.140$.	

Question 86

Skill	Learning Objective	Topic
2.B	FUN-4.A	Connecting a Function, Its First Derivative, and Its Second Derivative
(A)	Correct. The graph of f indicates that f is increasing from $x = 0$ to $x = 2$, then decreasing from $x = 2$ to $x = 3$, and then increasing from $x = 3$ to $x = 5$. Therefore, the graph of f' should be positive from $x = 0$ to $x = 2$, negative from $x = 2$ to $x = 3$, and positive from $x = 3$ to $x = 5$. This graph is the only one that has this behavior, so it could be the graph of f'. Some other features of the graph of f support this conclusion. Since f is not differentiable at $x = 2$, the graph of f' should not be defined at $x = 2$. Since f has a local minimum at $x = 3$ and is differentiable there, $f'(3)$ should equal 0. This graph is consistent with those observations.	
(B)	Incorrect. This graph shows the correct sign for f' on the interval from $x = 0$ to $x = 2$. However, the sign of f' should be negative from $x = 2$ to $x = 3$ and positive from $x = 3$ to $x = 5$, since f changes from decreasing to increasing over the interval $(2, 5)$. That is the opposite of what is happening between $x = 2$ and $x = 5$ in this graph. In addition, the graph of f does not support the conclusion that $\lim_{x \rightarrow 2^-} f'(x) = \lim_{x \rightarrow 2^+} f'(x)$, as suggested in this graph.	
(C)	Incorrect. This graph shows the correct sign for f' on the interval from $x = 2$ to $x = 5$. However, the sign of f' should be positive from $x = 0$ to $x = 2$, not negative, since f is increasing from $x = 0$ to $x = 2$. In addition, the graph of f does not support the conclusion that $\lim_{x \rightarrow 2^-} f'(x) = \lim_{x \rightarrow 2^+} f'(x)$, as suggested in this graph.	
(D)	Incorrect. This graph shows the correct sign for f' on the interval from $x = 0$ to $x = 3$. However, the sign of f' should be positive from $x = 3$ to $x = 5$, since f is increasing over the interval $(3, 5)$. In addition, the graph of f does not support the conclusion that $\lim_{x \rightarrow 2^+} f'(x) = 0$, as suggested in this graph.	

Question 87

Skill	Learning Objective	Topic
1.E	CHA-4.C	Connecting Position, Velocity, and Acceleration Functions Using Integrals
(A)	Incorrect. It was correctly determined that the particle changes direction from moving left to moving right at $t = b = 1.84527$. This value might have been stored in the calculator, and the stored value used for the limit of integration in order to ensure accuracy. However, this response is the displacement during the time interval $0 < t < b$, not the total distance traveled. $\int_0^b v(t) dt = \int_0^{1.84527} t \sin(t^3) dt = 0.212$	
(B)	Incorrect. The zero of $v(t)$ where the velocity changes from positive to negative was found to be $t = a = 1.46459$. This value might have been stored in the calculator, and the stored value used for the limit of integration in order to ensure accuracy. This is the time when the particle changes direction from moving right to moving left, not from left to right. The total distance traveled by the particle during the time interval $0 < t < a$ is $\int_0^a v(t) dt = \int_0^{1.46459} t \sin(t^3) dt = 0.612.$	
(C)	Correct. The graph of the velocity over the interval $(0, 2)$ shows that the velocity changes from positive to negative, then back to positive. The time at which the particle changes direction from moving left to moving right, therefore, is the second zero of $v(t)$, where the velocity changes from negative to positive. This zero is at $t = b = 1.84527$. Store this value in the calculator, and use the stored value for the limit of integration in order to ensure accuracy. The total distance traveled by the particle during the time interval $0 < t < b$ is $\int_0^b v(t) dt = \int_0^{1.84527} t \sin(t^3) dt = 1.011.$	
(D)	Incorrect. This response is the total distance traveled by the particle during the entire time interval $0 < t < 2$. $\int_0^2 v(t) dt = \int_0^2 t \sin(t^3) dt = 1.208$	

Question 88

Skill	Learning Objective	Topic
1.E	CHA-3.A	Interpreting the Meaning of the Derivative in Context
(A)	Incorrect. This response might be chosen if the calculation of the average rate of change resulted in a value that was greater than 0 or less than -0.5 . It would also be chosen if the average rate of change was correctly found to be -0.39206 , but the instantaneous rate of change was taken to be the second derivative of f , not the first derivative. In either case, the resulting equation would have no solution in the interval $[0, 1.565]$.	
(B)	Incorrect. This response would be chosen if the average rate of change was correctly found to be -0.39206 , but the graph of f , not f' , was drawn to determine the number of intersection points with the horizontal line $y = -0.39206$. It would also be chosen if the instantaneous rate of change was correctly identified as the derivative of f , but the average rate of change over the interval $[0, 1.565]$ was thought to be the average at the endpoints, $\frac{f(0) + f(1.565)}{2} = -0.30678$, or the average value of the function over the interval, $\frac{1}{1.565} \int_0^{1.565} f(x) dx = -0.32195$. In all these cases, the resulting equation would have only one solution in the interval $[0, 1.565]$.	
(C)	Correct. The average rate of change of f on the closed interval $[0, 1.565]$ is $\frac{f(1.565) - f(0)}{1.565 - 0} = -0.39206$. The instantaneous rate of change of f is the derivative, $f'(x) = x^3 - 2x^2 + x - \frac{1}{2}$. The graph of f' , produced using the calculator, intersects the horizontal line $y = -0.39206$ three times in the open interval $(0, 1.565)$.	
(D)	Incorrect. This response might be chosen because the function f is a polynomial of degree 4.	

Question 89

Skill	Learning Objective	Topic
3.E	FUN-4.A	Sketching Graphs of Functions and Their Derivatives
(A)	Incorrect. The graph of g is increasing because $g'(x) > 0$ and concave up because $g''(x) > 0$. The secant line through the points $(3, 12)$ and $(5, 18)$ is $y = 3(x - 3) + 12$. Because the graph of g is increasing and concave up, the graph will lie above the secant line for $x > 5$. In particular, the value of $g(6)$ is strictly greater than the value of y on the secant line at $x = 6$, that is, $g(6) > 3(6 - 3) + 12 = 21$. Therefore, $g(6)$ cannot equal 21.	
(B)	Correct. The graph of g is increasing because $g'(x) > 0$ and concave up because $g''(x) > 0$. The secant line through the points $(3, 12)$ and $(5, 18)$ is $y = 3(x - 3) + 12$. Because the graph of g is increasing and concave up, the graph will lie above the secant line for $x > 5$. In particular, the value of $g(6)$ is strictly greater than the value of y on the secant line at $x = 6$, that is, $g(6) > 3(6 - 3) + 12 = 21$. Therefore, 22 is the only possible value for $g(6)$.	
(C)	Incorrect. The graph of g is increasing because $g'(x) > 0$ and concave up because $g''(x) > 0$. The secant line through the points $(3, 12)$ and $(5, 18)$ is $y = 3(x - 3) + 12$. Because the graph of g is increasing and concave up, the graph will lie above the secant line for $x > 5$. In particular, the value of $g(6)$ is strictly greater than the value of y on the secant line at $x = 6$, that is, $g(6) > 3(6 - 3) + 12 = 21$. Therefore, $g(6)$ cannot equal 20 or 21.	
(D)	Incorrect. The graph of g is increasing because $g'(x) > 0$ and concave up because $g''(x) > 0$. The secant line through the points $(3, 12)$ and $(5, 18)$ is $y = 3(x - 3) + 12$. Because the graph of g is increasing and concave up, the graph will lie above the secant line for $x > 5$. In particular, the value of $g(6)$ is strictly greater than the value of y on the secant line at $x = 6$, that is, $g(6) > 3(6 - 3) + 12 = 21$. Therefore, $g(6)$ could equal 22, but not 21.	

Question 90

Skill	Learning Objective	Topic
1.E	FUN-3.E	Differentiating Inverse Functions
(A)	Incorrect. This response would result if it was correctly determined that $g(2) = f^{-1}(2) = 3$, but a sign error was made in taking $g'(2) = -\frac{1}{f'(3)}$ rather than $g'(2) = \frac{1}{f'(3)}$, perhaps by combining the calculations of the slope of an inverse function and the slope of a perpendicular line.	
(B)	Incorrect. This response would result if the derivative of g at $x = 2$ was taken to be the reciprocal of the derivative of f at $x = 2$, resulting in $g'(2) = \frac{1}{f'(2)} = \frac{1}{4}$ instead of $g'(2) = \frac{1}{f'(g(2))} = \frac{1}{f'(3)} = \frac{1}{5}$. In addition, the line was computed using the point $(2, 1)$ on the graph of f rather than the point $(2, 3)$ on the graph of g .	
(C)	Correct. Since $f(g(x)) = x$, the chain rule can be used to determine that $f'(g(x))g'(x) = 1$. Substituting $x = 2$ gives $1 = f'(g(2))g'(2) = f'(3)g'(2) \Rightarrow g'(2) = \frac{1}{f'(3)} = \frac{1}{5}$. Since $g(2) = f^{-1}(2) = 3$, an equation of the line tangent to the graph of g at $x = 2$ is therefore $y = \frac{1}{5}(x - 2) + 3$.	
(D)	Incorrect. This response is an equation of the line tangent to the graph of f at the point where $x = 2$ rather than the line tangent to the graph of g , the inverse function of f , at $x = 2$.	

AP[®] CALCULUS AB
2019 SCORING GUIDELINES

Question 1

(a) $E'(7) = 6.164924$

The rate of change of $E(t)$ at time $t = 7$ is 6.165 (or 6.164) cars per hour per hour.

1 : answer with units

(b) $\int_0^{12} E(t) dt = 520.070489$

To the nearest whole number, 520 cars enter the parking lot from time $t = 0$ to time $t = 12$.

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(c)
$$\begin{aligned} \int_2^{12} L(t) dt &\approx (5-2) \cdot \frac{L(2)+L(5)}{2} + (9-5) \cdot \frac{L(5)+L(9)}{2} \\ &\quad + (11-9) \cdot \frac{L(9)+L(11)}{2} + (12-11) \cdot \frac{L(11)+L(12)}{2} \\ &= 3 \cdot \frac{15+40}{2} + 4 \cdot \frac{40+24}{2} + 2 \cdot \frac{24+68}{2} + 1 \cdot \frac{68+18}{2} \\ &= 345.5 \end{aligned}$$

$\int_2^{12} L(t) dt$ is the number of cars that leave the parking lot in the 10 hours between 7 A.M. ($t = 2$) and 5 P.M. ($t = 12$).

3 : $\begin{cases} 1 : \text{trapezoidal sum} \\ 1 : \text{approximation} \\ 1 : \text{explanation} \end{cases}$

(d) $5 \int_0^6 E(t) dt + 8 \int_6^{12} E(t) dt = 3530.1396$

To the nearest dollar, 3530 dollars are collected from time $t = 0$ to time $t = 12$.

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{limits and constants} \\ 1 : \text{answer} \end{cases}$

AP[®] CALCULUS AB
2019 SCORING GUIDELINES

Question 2

$$\begin{aligned} \text{(a)} \quad \int_0^3 (f(x) - g(x)) \, dx &= \int_0^3 f(x) \, dx - \int_0^3 g(x) \, dx \\ &= \int_0^3 f(x) \, dx - 3.24125 = 4.919585 \end{aligned}$$

The area of region S is 4.920 (or 4.919).

3 : $\begin{cases} 1 : \text{definite integral of } f \\ 1 : \text{uses area of } R \\ 1 : \text{answer} \end{cases}$

$$\begin{aligned} \text{(b)} \quad \pi \int_0^3 ((f(x) + 3)^2 - (g(x) + 3)^2) \, dx \\ &= \pi \left(\int_0^3 (f(x) + 3)^2 \, dx - \int_0^3 ((g(x))^2 + 6g(x) + 9) \, dx \right) \\ &= \pi \left(\int_0^3 (f(x) + 3)^2 \, dx - \int_0^3 (g(x))^2 \, dx - 6 \int_0^3 g(x) \, dx - \int_0^3 9 \, dx \right) \\ &= \pi \left(\int_0^3 (f(x) + 3)^2 \, dx - 5.32021 - 6 \cdot 3.24125 - 9 \cdot 3 \right) \\ &= 156.263709 \end{aligned}$$

The volume of the solid is 156.264 (or 156.263).

4 : $\begin{cases} 1 : \text{form of integrand} \\ 1 : \text{integrand} \\ 1 : \text{uses areas of } R \text{ and } T \\ 1 : \text{limits, constant, and answer} \end{cases}$

$$\text{(c)} \quad \text{Volume} = \int_0^3 7(f(x) - g(x))^2 \, dx$$

2 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{expression} \end{cases}$

AP[®] CALCULUS AB
2019 SCORING GUIDELINES

Question 3

(a) Average rate of change = $\frac{f(4) - f(-3)}{4 - (-3)} = \frac{-1 - 0}{7} = -\frac{1}{7}$

1 : answer

(b) $f(3) = -3 + 3\cos\left(\frac{3\pi}{2}\right) = -3$

2 : $\begin{cases} 1 : f'(3) \\ 1 : \text{equation} \end{cases}$

For $0 < x < 4$, $f'(x) = -1 + \left(-3\sin\left(\frac{\pi x}{2}\right)\right) \cdot \frac{\pi}{2}$

$f'(3) = -1 + \left(-3\sin\left(\frac{3\pi}{2}\right)\right) \cdot \frac{\pi}{2} = -1 + \frac{3\pi}{2}$

An equation for the tangent line is $y = -3 + \left(-1 + \frac{3\pi}{2}\right)(x - 3)$.

(c) The average value of f on the interval $-3 \leq x \leq 4$ is

$\frac{1}{4 - (-3)} \int_{-3}^4 f(x) dx$.

4 : $\begin{cases} 1 : \text{integrals of } f \text{ over} \\ \quad -3 \leq x \leq 0 \text{ and } 0 \leq x \leq 4 \\ 1 : \text{value of } \int_{-3}^0 \sqrt{9 - x^2} dx \\ 1 : \text{antiderivative of} \\ \quad -x + 3\cos\left(\frac{\pi x}{2}\right) \\ 1 : \text{answer} \end{cases}$

$\int_{-3}^4 f(x) dx = \int_{-3}^0 f(x) dx + \int_0^4 f(x) dx$

$\int_{-3}^0 f(x) dx = \int_{-3}^0 \sqrt{9 - x^2} dx = \frac{9\pi}{4}$

$\int_0^4 f(x) dx = \int_0^4 \left(-x + 3\cos\left(\frac{\pi x}{2}\right)\right) dx = \left[-\frac{1}{2}x^2 + \frac{6}{\pi}\sin\left(\frac{\pi x}{2}\right)\right]_0^4 = -8$

$\frac{1}{4 - (-3)} \int_{-3}^4 f(x) dx = \frac{1}{7} \left(\frac{9\pi}{4} - 8\right)$

(d) $\lim_{x \rightarrow 0^-} f(x) = f(0) = 3$ and $\lim_{x \rightarrow 0^+} f(x) = 3$, so f is continuous at $x = 0$.

2 : $\begin{cases} 1 : \text{continuity at } x = 0 \\ 1 : \text{answer with justification} \end{cases}$

Because f is continuous on $[-3, 4]$, the Extreme Value Theorem guarantees that f attains an absolute maximum on $[-3, 4]$.

AP[®] CALCULUS AB
2019 SCORING GUIDELINES

Question 4

(a) $g(0) = \int_{-4}^0 f(t) \, dt = \frac{9}{2} - 3 = \frac{3}{2}$

$$\begin{aligned} g(4) &= \int_{-4}^4 f(t) \, dt \\ &= \int_{-4}^0 f(t) \, dt + \int_0^1 f(t) \, dt + \int_1^4 f(t) \, dt \\ &= \frac{3}{2} + 5 + \int_1^4 (-t^2 + 5t - 4) \, dt \\ &= \frac{3}{2} + 5 + \left[-\frac{1}{3}t^3 + \frac{5}{2}t^2 - 4t \right]_1^4 \\ &= \frac{3}{2} + 5 + \left[\left(-\frac{1}{3} \cdot 4^3 + \frac{5}{2} \cdot 4^2 - 4 \cdot 4 \right) - \left(-\frac{1}{3} \cdot 1^3 + \frac{5}{2} \cdot 1^2 - 4 \cdot 1 \right) \right] \\ &= \frac{3}{2} + 5 + \left(\frac{8}{3} - \left(-\frac{11}{6} \right) \right) = 11 \end{aligned}$$

- (b) $g'(x) = f(x)$ is negative for $-1 < x < 0$, and nonnegative elsewhere. Thus, the absolute minimum value of g on $[-4, 4]$ can only occur at $x = -4$ or $x = 0$.

$$g(-4) = 0$$

$$g(0) = \frac{3}{2}$$

The absolute minimum value of g on $[-4, 4]$ is $g(-4) = 0$.

- (c) The graph of g is concave down on the intervals $-2 < x < -\frac{1}{2}$, $\frac{1}{2} < x < 1$, and $\frac{5}{2} < x < 4$ because $g'(x) = f(x)$ is decreasing on these intervals.

$$4 : \begin{cases} 1 : g(0) \\ 1 : \text{integral of } f \text{ over } 1 \leq t \leq 4 \\ 1 : \text{antiderivative} \\ 1 : g(4) \end{cases}$$

$$3 : \begin{cases} 1 : g'(x) = f(x) \\ 1 : \text{identifies } x = -4 \text{ and } x = 0 \\ \quad \text{as candidates} \\ 1 : \text{answer with justification} \end{cases}$$

$$2 : \begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$$

AP[®] CALCULUS AB
2019 SCORING GUIDELINES

Question 5

(a) $\int_0^2 te^{4-t^2} dt = -\frac{1}{2}e^{4-t^2} \Big|_{t=0}^{t=2} = -\frac{1}{2} + \frac{1}{2}e^4$

Chloe traveled $-\frac{1}{2} + \frac{1}{2}e^4$ miles from time $t = 0$ to time $t = 2$.

$$3 : \begin{cases} 1 : \text{integral} \\ 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$$

(b) $C(3) = -6 < 0$
 $C'(3) = -9 < 0$

Chloe's speed is increasing at time $t = 3$ because her velocity and acceleration have the same sign.

$$2 : \begin{cases} 1 : C(3) < 0 \text{ and } C'(3) < 0 \\ 1 : \text{answer with reason} \end{cases}$$

(c) B is differentiable $\Rightarrow B$ is continuous on $[0, 4]$.

$$\frac{B(4) - B(0)}{4 - 0} = \frac{11 - 1}{4 - 0} = 2.5$$

By the Mean Value Theorem, there is a time t , for $0 < t < 4$, such that $B'(t) = 2.5$ miles per hour per hour.

$$2 : \begin{cases} 1 : \frac{B(4) - B(0)}{4 - 0} \\ 1 : \text{answer with justification} \end{cases}$$

(d) B and C are continuous on $[0, 2]$, therefore $B - C$ is continuous on $[0, 2]$.

$$B(0) - C(0) = 1 - 0 > 0$$

$$B(2) - C(2) = 1.5 - 2 < 0$$

By the Intermediate Value Theorem, there is a time t , for $0 < t < 2$, such that $B(t) - C(t) = 0$, or $B(t) = C(t)$.

$$2 : \begin{cases} 1 : \text{considers } B(t) - C(t) \\ 1 : \text{answer with justification} \end{cases}$$

AP[®] CALCULUS AB
2019 SCORING GUIDELINES

Question 6

(a) $\frac{d}{dx}(2x^2 + 3y^2 - 4xy) = \frac{d}{dx}(36) \Rightarrow 4x + 6y \frac{dy}{dx} - 4y - 4x \frac{dy}{dx} = 0$
 $\Rightarrow (6y - 4x) \frac{dy}{dx} = 4y - 4x \Rightarrow \frac{dy}{dx} = \frac{4y - 4x}{6y - 4x} = \frac{2y - 2x}{3y - 2x}$

2 : $\begin{cases} 1 : \text{implicit differentiation} \\ 1 : \text{verification} \end{cases}$

(b) $x = 6 \Rightarrow 2 \cdot 6^2 + 3y^2 - 4 \cdot 6 \cdot y = 36$
 $\Rightarrow y^2 - 8y + 12 = 0 \Rightarrow y = 2 \text{ or } y = 6$
 $\left. \frac{dy}{dx} \right|_{(x,y)=(6,2)} = \frac{4 - 12}{6 - 12} = \frac{4}{3}$

2 : $\begin{cases} 1 : \text{slope at } (6, 2) \\ 1 : \text{slope at } (6, 6) \end{cases}$

The slope of the line tangent to the curve at $(6, 2)$ is $\frac{4}{3}$.

$$\left. \frac{dy}{dx} \right|_{(x,y)=(6,6)} = \frac{12 - 12}{18 - 12} = 0$$

The slope of the line tangent to the curve at $(6, 6)$ is 0.

(c) The curve has vertical tangent lines where $3y - 2x = 0$ and $2y - 2x \neq 0$.

2 : $\begin{cases} 1 : \text{sets } 3y - 2x = 0 \\ 1 : \text{answer} \end{cases}$

$$3y - 2x = 0 \Rightarrow y = \frac{2}{3}x \Rightarrow 2x^2 + 3\left(\frac{2}{3}x\right)^2 - 4x \cdot \frac{2}{3}x = 36$$

$$\Rightarrow \left(2 + \frac{4}{3} - \frac{8}{3}\right)x^2 = \frac{2}{3}x^2 = 36 \Rightarrow x^2 = 54$$

Because x is positive, $x = 3\sqrt{6}$.

(d) $\frac{d}{dt}(2x^2 + 3y^2 - 4xy) = \frac{d}{dt}(36)$
 $\Rightarrow 4x \frac{dx}{dt} + 6y \frac{dy}{dt} - 4\left(x \frac{dy}{dt} + y \frac{dx}{dt}\right) = 0$
 $\Rightarrow (4x - 4y) \frac{dx}{dt} + (6y - 4x) \frac{dy}{dt} = 0$

3 : $\begin{cases} 1 : \text{chain rules or product rule} \\ 1 : \text{derivative with respect to } t \\ 1 : \text{answer} \end{cases}$

$$(4 \cdot 2 - 4 \cdot (-2)) \frac{dx}{dt} \Big|_{t=1} + (6 \cdot (-2) - 4 \cdot 2) \cdot 4 = 0 \Rightarrow \frac{dx}{dt} \Big|_{t=1} = 5$$

— OR —

$$\frac{dy}{dt} = \frac{dy}{dx} \cdot \frac{dx}{dt} = \frac{2y - 2x}{3y - 2x} \cdot \frac{dx}{dt}$$

$$4 = \frac{2 \cdot (-2) - 2 \cdot 2}{3 \cdot (-2) - 2 \cdot 2} \cdot \frac{dx}{dt} \Big|_{t=1} = \frac{4}{5} \cdot \frac{dx}{dt} \Big|_{t=1} \Rightarrow \frac{dx}{dt} \Big|_{t=1} = 5$$

2019 AP Calculus AB Scoring Worksheet

Section I: Multiple Choice

$$\frac{\text{Number Correct (out of 45)}}{\text{Number Correct (out of 45)}} \times 1.2000 = \frac{\text{Weighted Section I Score (Do not round)}}{\text{Weighted Section I Score (Do not round)}}$$

Section II: Free Response

$$\frac{\text{Question 1 (out of 9)}}{\text{Question 1 (out of 9)}} \times 1.0000 = \frac{\text{(Do not round)}}{\text{(Do not round)}}$$

$$\frac{\text{Question 2 (out of 9)}}{\text{Question 2 (out of 9)}} \times 1.0000 = \frac{\text{(Do not round)}}{\text{(Do not round)}}$$

$$\frac{\text{Question 3 (out of 9)}}{\text{Question 3 (out of 9)}} \times 1.0000 = \frac{\text{(Do not round)}}{\text{(Do not round)}}$$

$$\frac{\text{Question 4 (out of 9)}}{\text{Question 4 (out of 9)}} \times 1.0000 = \frac{\text{(Do not round)}}{\text{(Do not round)}}$$

$$\frac{\text{Question 5 (out of 9)}}{\text{Question 5 (out of 9)}} \times 1.0000 = \frac{\text{(Do not round)}}{\text{(Do not round)}}$$

$$\frac{\text{Question 6 (out of 9)}}{\text{Question 6 (out of 9)}} \times 1.0000 = \frac{\text{(Do not round)}}{\text{(Do not round)}}$$

$$\text{Sum} = \frac{\text{Weighted Section II Score (Do not round)}}{\text{Weighted Section II Score (Do not round)}}$$

Composite Score

$$\frac{\text{Weighted Section I Score}}{\text{Weighted Section I Score}} + \frac{\text{Weighted Section II Score}}{\text{Weighted Section II Score}} = \frac{\text{Composite Score (Round to nearest whole number)}}{\text{Composite Score (Round to nearest whole number)}}$$

AP Score Conversion Chart
Calculus AB

Composite Score Range	AP Score
69-108	5
56-68	4
43-55	3
27-42	2
0-26	1

2019 AP Calculus AB

Question Descriptors and Performance Data

Multiple-Choice Questions

Question	Skill	Learning Objective	Topic	Key	% Correct
1	1.E	FUN-6.C	Finding Antiderivatives and Indefinite Integrals - Basic Rules and Notation	B	88
2	1.D	CHA-2.C	Defining the Derivative of a Function and Using Derivative Notation	C	65
3	1.E	FUN-3.C	The Chain Rule	D	66
4	1.E	FUN-6.D	Integrating Using Substitution	D	54
5	2.B	FUN-4.A	Determining Concavity of Functions over Their Domains	C	70
6	1.E	FUN-3.D	Implicit Differentiation	D	41
7	1.E	FUN-3.A	Derivatives of $\cos x$, $\sin x$, e^x , and $\ln x$	D	62
8	2.B	LIM-2.D	Connecting Limits at Infinity and Horizontal Asymptotes	B	50
9	1.E	CHA-5.A	Finding the Area Between Curves Expressed as Functions of x	B	64
10	1.E	FUN-3.B	The Quotient Rule	A	64
11	1.E	FUN-6.A	Applying Properties of Definite Integrals	A	53
12	3.D	FUN-1.B	Using the Mean Value Theorem	C	63
13	1.E	LIM-4.A	Using L'Hospital's Rule for Finding Limits of Indeterminate Forms	A	60
14	2.C	FUN-7.F	Exponential Models with Differential Equations	D	75
15	3.C	FUN-2.A	Connecting Differentiability and Continuity - Determining When Derivatives Do and Do Not Exist	A	54
16	1.E	FUN-7.D	Finding General Solutions Using Separation of Variables	B	30
17	2.B	FUN-5.A	Interpreting the Behavior of Accumulation Functions Involving Area	B	65
18	1.E	FUN-3.B	The Product Rule	D	62
19	2.B	CHA-3.B	Straight-Line Motion: Connecting Position, Velocity, and Acceleration	A	34
20	1.C	FUN-6.A	Applying Properties of Definite Integrals	C	57
21	1.D	CHA-2.B	Defining the Derivative of a Function and Using Derivative Notation	D	61
22	2.D	FUN-7.C	Sketching Slope Fields	D	49
23	1.E	CHA-5.B	Volumes with Cross Sections - Squares and Rectangles	B	36
24	3.D	FUN-6.B	The Fundamental Theorem of Calculus and Definite Integrals	A	40
25	3.G	FUN-7.B	Verifying Solutions for Differential Equations	A	34
26	3.D	FUN-4.A	Using the Candidates Test to Find Absolute (Global) Extrema	B	53
27	3.F	CHA-3.A	Interpreting the Meaning of the Derivative in Context	C	66
28	1.E	CHA-3.E	Solving Related Rates Problems	B	36
29	2.B	LIM-2.D	Connecting Infinite Limits and Vertical Asymptotes	A	34
30	3.D	LIM-5.C	Riemann Sums, Summation Notation, and Definite Integral Notation	B	27

2019 AP Calculus AB

Question Descriptors and Performance Data

Question	Skill	Learning Objective	Topic	Key	% Correct
76	2.E	FUN-4.A	Determining Intervals on Which a Function Is Increasing or Decreasing	D	82
77	1.E	CHA-3.B	Straight-Line Motion: Connecting Position, Velocity, and Acceleration	B	86
78	1.E	CHA-3.F	Approximating Values of a Function Using Local Linearity and Linearization	A	66
79	3.F	CHA-4.B	Finding the Average Value of a Function on an Interval	B	61
80	1.E	FUN-6.B	The Fundamental Theorem of Calculus and Definite Integrals	C	62
81	2.B	LIM-2.A	Exploring Types of Discontinuities	C	81
82	1.E	FUN-6.A	Applying Properties of Definite Integrals	A	71
83	3.D	LIM-2.C	Removing Discontinuities	A	53
84	3.D	FUN-1.A	Working with the Intermediate Value Theorem	C	48
85	3.D	FUN-4.A	Using the First Derivative Test to Find Relative (Local) Extrema	D	51
86	2.B	FUN-4.A	Connecting a Function, Its First Derivative, and Its Second Derivative	A	76
87	1.E	CHA-4.C	Connecting Position, Velocity, and Acceleration Functions Using Integrals	C	32
88	1.E	CHA-3.A	Interpreting the Meaning of the Derivative in Context	C	33
89	3.E	FUN-4.A	Sketching Graphs of Functions and Their Derivatives	B	33
90	1.E	FUN-3.E	Differentiating Inverse Functions	C	37

Free-Response Questions

Question	Skill	Learning Objective	Topic	Mean Score
1	1.E 3.D 3.F 4.D 4.A 4.B 4.C 4.E	CHA-2.D CHA-4.E LIM-5.A	2.3 8.3 6.2 8.3	6.03
2	1.D 1.E 2.B 4.E	CHA-5.A CHA-5.C CHA-5.B	8.4 8.12 8.7	3.46
3	1.D 1.E 3.C 3.D 4.A	CHA-2.A CHA-2.C CHA-4.B FUN-1.C	2.1 2.2 8.1 5.2	1.71
4	1.D 1.E 2.B 2.E 4.A	FUN-6.A FUN-4.A	6.6 5.5 5.6	2.98
5	1.D 1.E 3.D 3.E 4.A	CHA-4.C CHA-3.B FUN-1.B FUN-1.A	8.2 4.2 5.1 1.16	2.71
6	1.C 1.E 3.G 4.C	FUN-3.D FUN-4.E CHA-3.E	3.2 5.12 4.5	2.98

AP Calculus AB

The College Board

The College Board is a mission-driven not-for-profit organization that connects students to college success and opportunity. Founded in 1900, the College Board was created to expand access to higher education. Today, the membership association is made up of over 6,000 of the world's leading educational institutions and is dedicated to promoting excellence and equity in education. Each year, the College Board helps more than seven million students prepare for a successful transition to college through programs and services in college readiness and college success — including the SAT[®] and the Advanced Placement Program[®]. The organization also serves the education community through research and advocacy on behalf of students, educators, and schools. The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities, and concerns.