
Please note: Some of the questions in this former practice exam may no longer perfectly align with the AP exam. Even though these questions do not fully represent the 2020 exam, teachers indicate that imperfectly aligned questions still provide instructional value. Teachers can consult the Question Bank to determine the degree to which these questions align to the 2020 Exam.

This exam may not be posted on school or personal websites, nor electronically redistributed for any reason. This exam is provided by the College Board for AP Exam preparation. Teachers are permitted to download the materials and make copies to use with their students in a classroom setting only. To maintain the security of this exam, teachers should collect all materials after their administration and keep them in a secure location.

Further distribution of these materials outside of the secure College Board site disadvantages teachers who rely on uncirculated questions for classroom testing. Any additional distribution is in violation of the College Board's copyright policies and may result in the termination of Practice Exam access for your school as well as the removal of access to other online services such as the AP Teacher Community and Online Score Reports.

AP[®] Chemistry

Practice Exam

NOTE: This is a modified version of the 2019 AP Chemistry Exam.

This exam may not be posted on school or personal websites, nor electronically redistributed for any reason. This Released Exam is provided by the College Board for AP Exam preparation. Teachers are permitted to download the materials and make copies to use with their students in a classroom setting only. To maintain the security of this exam, teachers should collect all materials after their administration and keep them in a secure location.

Further distribution of these materials outside of the secure College Board site disadvantages teachers who rely on uncirculated questions for classroom testing. Any additional distribution is in violation of the College Board's copyright policies and may result in the termination of Practice Exam access for your school as well as the removal of access to other online services such as the AP Teacher Community and Online Score Reports.

Contents

Exam Instructions

Student Answer Sheet for the Multiple-Choice Section

Section I: Multiple-Choice Questions

Section II: Free-Response Questions

Multiple-Choice Answer Key

Course Framework Alignment and Rationales

Free-Response Scoring Guidelines

Scoring Worksheet

Question Descriptors and Performance Data

Note: This publication shows the page numbers that appeared in the **2018–19 AP Exam Instructions** book and in the actual exam. This publication was not repaginated to begin with page 1.

AP Chemistry Exam

Regularly Scheduled Exam Date: Thursday morning, May 9, 2019

Late-Testing Exam Date: Wednesday afternoon, May 22, 2019

Section I	Total Time: 1 hour and 30 minutes Number of Questions: 50 <i>(The number of questions may vary slightly depending on the form of the exam.)</i> Calculator not permitted Percent of Total Score: 50% Writing Instrument: Pencil required
Section II	Total Time: 1 hour and 45 minutes Number of Questions: 7 <i>(3 ten-point and 4 four-point questions)</i> Calculators allowed for all of Section II Percent of Total Score: 50% Writing Instrument: Either pencil or pen with black or dark blue ink

Before Distributing Exams: Check that the title on all exam covers is **Chemistry**. If there are any exam booklets with a different title, contact the AP coordinator immediately.

Note: Tables of equations and constants are provided in the exam booklets for both sections of the exam.

What Proctors Need to Bring to This Exam

- ☐ Exam packets
- ☐ Answer sheets
- ☐ AP Student Packs
- ☐ 2018-19 AP Coordinator's Manual
- ☐ This book—2018-19 AP Exam Instructions
- ☐ AP Exam Seating Chart template
- ☐ School Code and Homeschool/Self-Study Codes
- ☐ Extra calculators
- ☐ Pencil sharpener
- ☐ Container for students' electronic devices (if needed)
- ☐ Extra No. 2 pencils with erasers
- ☐ Extra pens with black or dark blue ink
- ☐ Extra paper
- ☐ Stapler
- ☐ Watch
- ☐ Signs for the door to the testing room
 - “Exam in Progress”
 - “Phones of any kind are prohibited during the test administration, including breaks”

Students are not allowed to use calculators in Section I of the AP Chemistry Exam. However, students are permitted to use scientific or graphing calculators to answer questions in Section II. Four-function calculators are also permitted for use in Section II, but are not recommended. Before starting the exam administration, make sure that each student has an appropriate calculator and that any student with a graphing calculator has a model from the approved list on page 53 of the *2018-19 AP Coordinator's Manual*. See pages 50–53 of the *AP Coordinator's Manual* for more information. If a student does not have an appropriate calculator or has a graphing calculator not on the approved list, you may provide one from your supply. If the student does not want to use the calculator you provide, or does not want to use a calculator at all, he or she must hand copy, date, and sign the release statement on page 52 of the *AP Coordinator's Manual*.

During the administration of Section II, students may have no more than two calculators on their desks. Calculators may not be shared. Calculator memories do not need to be cleared before or after the exam. Students with Hewlett-Packard 48–50 Series and Casio FX-9860 graphing calculators may use cards designed for use with these calculators. Proctors should make sure infrared ports (Hewlett-Packard) are not facing each other. **Since graphing calculators can be used to store data, including text, proctors should monitor that students are using their calculators appropriately. Attempts by students to use the calculator to remove exam questions and/or answers from the room may result in the cancellation of AP Exam scores.**

Students will be allowed to use the table of equations and constants on both sections of the exam.

NOTE: The Periodic Table of the Elements and the AP Chemistry Equations and Constants must not be removed from the exam books.

SECTION I: Multiple Choice

► **Do not begin the exam instructions below until you have completed the appropriate General Instructions for your group.**

Make sure you begin the exam at the designated time. Remember, you must complete a seating chart for this exam. See pages 295–296 for a seating chart template and instructions. See the *2018-19 AP Coordinator's Manual* for exam seating requirements (pages 56–59).

If you are giving the regularly scheduled exam, say:

It is Thursday morning, May 9, and you will be taking the AP Chemistry Exam.

If you are giving the alternate exam for late testing, say:

It is Wednesday afternoon, May 22, and you will be taking the AP Chemistry Exam.

Look at your exam packet and confirm that the exam title is “AP Chemistry.”

Raise your hand if your exam packet contains any title other than “AP Chemistry,” and I will help you.

Once you confirm that all students have the correct exam, say:

In a moment, you will open the exam packet. By opening this packet, you agree to all of the AP Program's policies and procedures outlined in the *2018-19 Bulletin for AP Students and Parents*.

You may now remove the shrinkwrap from the outside only of your exam packet. Do not open the Section I booklet; do not remove the shrinkwrap from the Section II materials. Put the white seals and the shrinkwrapped Section II booklet aside. . . .

Carefully remove the AP Exam label found near the top left of your exam booklet cover. Place it on page 1 of your answer sheet on the light blue box near the top right corner that reads “AP Exam Label.”

If students accidentally place the exam label in the space for the number label or vice versa, advise them to leave the labels in place. They should not try to remove the label; their exam can still be processed correctly.

Listen carefully to all my instructions. I will give you time to complete each step. Please look up after completing each step. Raise your hand if you have any questions.

Give students enough time to complete each step. Don't move on until all students are ready.

Read the statements on the front cover of the Section I booklet. . . .

Sign your name and write today's date. . . .

Now print your full legal name where indicated. . . .

Turn to the back cover of your exam booklet and read it completely. . . .

Give students a few minutes to read the entire cover.

Are there any questions? . . .

You will now take the multiple-choice portion of the exam. You should have in front of you the multiple-choice booklet and your answer sheet. You may never discuss the multiple-choice exam content at any time in any form with anyone, including your teacher and other students. If you disclose the multiple-choice exam content through any means, your AP Exam score will be canceled.

Open your answer sheet to page 2. The answer sheet has circles A–E for each question. For Chemistry, you will use only the circles marked A–D. You must complete the answer sheet using a No. 2 pencil only. Mark all of your responses beginning on page 2 of your answer sheet, one response per question. Completely fill in the circles. If you need to erase, do so carefully and completely. No credit will be given for anything written in the exam booklet. Scratch paper is not allowed, but you may use the margins or any blank space in the exam booklet for scratch work. Calculators are not allowed for this section. Please put your calculators under your chair. Are there any questions? . . .

You have 1 hour and 30 minutes for this section. Open your Section I booklet and begin.



Note Start Time _____ . Note Stop Time _____ .

Check that students are marking their answers in pencil on their answer sheets and that they have not opened their shrinkwrapped Section II booklets.

After 1 hour and 20 minutes, say:

There are 10 minutes remaining.

After 10 minutes, say:

Stop working. Close your booklet and put your answer sheet on your desk, faceup. Make sure you have your AP number label and an AP Exam label on page 1 of your answer sheet. Sit quietly while I collect your answer sheets.

Collect an answer sheet from each student. Check that each answer sheet has an AP number label and an AP Exam label.

After all answer sheets have been collected, say:

Now you must seal your exam booklet using the white seals you set aside earlier. Remove the white seals from the backing and press one on each area of your exam booklet cover marked "PLACE SEAL HERE." Fold each seal over the back cover. When you have finished, place the booklet on your desk, faceup.

I will now collect your Section I booklet. . . .

Collect a Section I booklet from each student. Check that each student has signed the front cover of the sealed Section I booklet.

There is a 10-minute break between Sections I and II.

When all Section I materials have been collected and accounted for and you are ready for the break, say:

Please listen carefully to these instructions before we take a 10-minute break. All items you placed under your chair at the beginning of this exam, including your Student Pack, must stay there, and you are not permitted to open or access them in any way. Leave your shrinkwrapped Section II packet on top of your desk during the break. You are not allowed to consult teachers, other students, notes, textbooks, or any other resources during the break. You may not make phone calls, send text messages, check email, use a social networking site, or access any electronic or communication device. You may not leave the designated break area. Remember, you may never discuss the multiple-choice exam content with anyone, and if you disclose the content through any means, your AP Exam score will be canceled. Are there any questions? . . .



You may begin your break. Testing will resume at _____ .

SECTION II: Free Response

After the break, say:

May I have everyone's attention? Place your Student Pack on your desk. . . .

You may now remove the shrinkwrap from the Section II packet, but do not open the exam booklet until you are told to do so. . . .

Read the bulleted statements on the front cover of the exam booklet. Look up when you have finished. . . .

Now take an AP number label from your Student Pack and place it on the shaded box. If you don't have any AP number labels, write your AP number in the box. Look up when you have finished. . . .

Read the last statement. . . .

Using your pen, print the first, middle, and last initials of your legal name in the boxes and print today's date where indicated. This constitutes your signature and your agreement to the statements on the front cover. . . .

Now turn to the back cover. Using your pen, complete Items 1 through 3 under "Important Identification Information." . . .

Read Item 4. . . .

Are there any questions? . . .

If this is your last AP Exam, you may keep your Student Pack. Place it under your chair for now. Otherwise if you are taking any other AP Exams this year, leave your Student Pack on your desk and I will collect it now. . . .

Read the remaining information on the back cover of the exam booklet. Do not open the exam booklet until you are told to do so. Look up when you have finished. . . .

Collect the Student Packs from students who are taking any other AP Exams this year.

Then say:

Are there any questions? . . .

Calculators may be used for Section II. You may get your calculators from under your chair and place them on your desk. . . .

You have 1 hour and 45 minutes to complete Section II. You are responsible for pacing yourself, and you may proceed freely from one question to the next. You must write your answers in the exam booklet using a pen with black or dark blue ink or a No. 2 pencil. If you use a pencil, be sure that your writing is dark enough to be easily read. If you need more paper to complete your responses, raise your hand. At the top of each extra sheet of paper you use, write only:

- your AP number,
- the exam title, and
- the question number you are working on.

Do not write your name. Are there any questions? . . .

You may begin.



Note Start Time _____ . Note Stop Time _____ .

Proctors should also make sure that Hewlett-Packard calculators' infrared ports are not facing each other and that students are not sharing calculators.

After 1 hour and 35 minutes, say:

There are 10 minutes remaining.

After 10 minutes, say:

Stop working and close your exam booklet. Place it on your desk, faceup. . . .

If any students used extra paper for a question in the free-response section, have those students staple the extra sheet(s) to the first page corresponding to that question in their free-response exam booklets. Complete an Incident Report after the exam and return these free-response booklets with the extra sheets attached in the Incident Report return envelope (see page 68 of the *2018-19 AP Coordinator's Manual* for complete details).

Then say:

Remain in your seat, without talking, while the exam materials are collected. . . .

Collect a Section II booklet from each student. Check for the following:

- Exam booklet front cover: The student placed an AP number label on the shaded box and printed their initials and today's date.
- Exam booklet back cover: The student completed the "Important Identification Information" area.

When all exam materials have been collected and accounted for, return to students any electronic devices you may have collected before the start of the exam.

If you are giving the regularly scheduled exam, say:

You may not discuss or share the free-response exam content with anyone unless it is released on the College Board website in about two days. Your AP Exam score results will be available online in July.

If you are giving the alternate exam for late testing, say:

None of the content in this exam may ever be discussed or shared in any way at any time. Your AP Exam score results will be available online in July.

If any students completed the AP number card at the beginning of this exam, say:

Please remember to take your AP number card with you. You will need the information on this card to view your scores and order AP score reporting services online.

Then say:

You are now dismissed.

After-Exam Tasks

Be sure to give the completed seating chart to the AP coordinator. Schools must retain seating charts for at least six months (unless the state or district requires that they be retained for a longer period of time). Schools should not return any seating charts in their exam shipments unless they are required as part of an Incident Report.

NOTE: If you administered exams to students with accommodations, review the *2018-19 AP Coordinator's Manual* and the *2018-19 AP SSD Guidelines* for information about completing the Nonstandard Administration Report (NAR) form, and returning these exams.

The exam proctor should complete the following tasks if asked to do so by the AP coordinator. Otherwise, the AP coordinator must complete these tasks:

- Complete an Incident Report for any students who used extra paper for the free-response section. (Incident Report forms are provided in the coordinator packets sent with the exam shipments.) **These forms must be completed with a No. 2 pencil.** It is best to complete a single Incident Report for multiple students per exam subject, per administration (regular or late testing), as long as all required information is provided. Include all exam booklets with extra sheets of paper in an Incident Report return envelope (see page 68 of the *2018-19 AP Coordinator's Manual* for complete details).
- Return all exam materials to secure storage until they are shipped back to the AP Program. (See page 27 of the *2018-19 AP Coordinator's Manual* for more information about secure storage.) Before storing materials, check the "School Use Only" section on page 1 of the answer sheet and:
 - ♦ Fill in the appropriate section number circle in order to access a separate AP Instructional Planning Report (for regularly scheduled exams only) or subject score roster at the class section or teacher level. See "Post-Exam Activities" in the *2018-19 AP Coordinator's Manual*.
 - ♦ Check your list of students who are eligible for fee reductions and fill in the appropriate circle on their registration answer sheets.

Name: _____

Answer Sheet for AP Chemistry
Practice Exam, Section I

No.	Answer
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	

No.	Answer
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	

AP[®] Chemistry Exam

SECTION I: Multiple Choice

2019

DO NOT OPEN THIS BOOKLET UNTIL YOU ARE TOLD TO DO SO.

At a Glance

Total Time

1 hour and 30 minutes

Number of Questions

50

Percent of Total Score

50%

Writing Instrument

Pencil required

Electronic Device

None allowed

Instructions

Section I of this exam contains 50 multiple-choice questions. Fill in only the circles for numbers 1 through 50 on your answer sheet. Pages containing a periodic table and lists containing equations and constants are also printed in this booklet.

Indicate all of your answers to the multiple-choice questions on the answer sheet. No credit will be given for anything written in this exam booklet, but you may use the booklet for notes or scratch work. After you have decided which of the suggested answers is best, completely fill in the corresponding circle on the answer sheet.

Because this section offers only four answer options for each question, do not mark the (E) answer circle for any question. Give only one answer to each question. If you change an answer, be sure that the previous mark is erased completely. Here is a sample question and answer.

Sample Question Sample Answer

Chicago is a

(A) ☒ (B) ☐ (C) ☐ (D) ☐ (E) ☐

(A) state

(B) city

(C) country

(D) continent

Use your time effectively, working as quickly as you can without losing accuracy. Do not spend too much time on any one question. Go on to other questions and come back to the ones you have not answered if you have time. It is not expected that everyone will know the answers to all of the multiple-choice questions.

Your total score on Section I is based only on the number of questions answered correctly. Points are not deducted for incorrect answers or unanswered questions.

Form I
Form Code 4BPB4-S

25

18

PERIODIC TABLE OF THE ELEMENTS

1

PERIODIC TABLE OF THE ELEMENTS																										
1		2										13					14		15		16		17		18	
H 1.008												5		6		7		8		9		10				
3 Li 6.94		4 Be 9.01										B 10.81		C 12.01		N 14.01		O 16.00		F 19.00		Ne 20.18				
11 Na 22.99		12 Mg 24.30										13 Al 26.98		14 Si 28.09		15 P 30.97		16 S 32.06		17 Cl 35.45		18 Ar 39.95				
19 K 39.10		20 Ca 40.08		3 21 Sc 44.96		4 22 Ti 47.87		5 23 V 50.94		6 24 Cr 52.00		7 25 Mn 54.94		8 26 Fe 55.85		9 27 Co 58.93		10 28 Ni 58.69		11 29 Cu 63.55		12 30 Zn 65.38		36 Kr 83.80		
37 Rb 85.47		38 Sr 87.62		39 Y 88.91		40 Zr 91.22		41 Nb 92.91		42 Mo 95.95		43 Tc (97)		44 Ru 101.1		45 Rh 102.91		46 Pd 106.42		47 Ag 107.87		48 Cd 112.41		54 Xe 131.29		
55 Cs 132.91		56 Ba 137.33		57 *La 138.91		72 Hf 178.49		73 Ta 180.95		74 W 183.84		75 Re 186.21		76 Os 190.2		77 Ir 192.2		78 Pt 195.08		79 Au 196.97		80 Hg 200.59		86 Rn (222)		
87 Fr (223)		88 Ra (226)		89 †Ac (227)		104 Rf (267)		105 Db (270)		106 Sg (271)		107 Bh (270)		108 Hs (277)		109 Mt (276)		110 Ds (281)		111 Rg (282)		112 Cn (285)		118 Uuo (294)		

*Lanthanoid Series

58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu
90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr

†Actinoid Series

AP[®] CHEMISTRY EQUATIONS AND CONSTANTS

Throughout the exam the following symbols have the definitions specified unless otherwise noted.

L, mL = liter(s), milliliter(s)
g = gram(s)
nm = nanometer(s)
atm = atmosphere(s)

mm Hg = millimeters of mercury
J, kJ = joule(s), kilojoule(s)
V = volt(s)
mol = mole(s)

ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

E = energy
 ν = frequency
 λ = wavelength

Planck's constant, $h = 6.626 \times 10^{-34}$ J s

Speed of light, $c = 2.998 \times 10^8$ m s⁻¹

Avogadro's number = 6.022×10^{23} mol⁻¹

Electron charge, $e = -1.602 \times 10^{-19}$ coulomb

EQUILIBRIUM

$$K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}, \text{ where } aA + bB \rightleftharpoons cC + dD$$

$$K_p = \frac{(P_C)^c(P_D)^d}{(P_A)^a(P_B)^b}$$

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[H^+], \text{ pOH} = -\log[OH^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[A^-]}{[HA]}$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

Equilibrium Constants

K_c (molar concentrations)

K_p (gas pressures)

K_a (weak acid)

K_b (weak base)

K_w (water)

KINETICS

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

k = rate constant

t = time

$t_{1/2}$ = half-life

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = ^\circ\text{C} + 273$$

$$D = \frac{m}{V}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

Molarity, M = moles of solute per liter of solution

$$A = abc$$

P = pressure

V = volume

T = temperature

n = number of moles

m = mass

M = molar mass

D = density

KE = kinetic energy

v = velocity

A = absorbance

a = molar absorptivity

b = path length

c = concentration

$$\begin{aligned}\text{Gas constant, } R &= 8.314 \text{ J mol}^{-1} \text{ K}^{-1} \\ &= 0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1} \\ &= 62.36 \text{ L torr mol}^{-1} \text{ K}^{-1}\end{aligned}$$

$$1 \text{ atm} = 760 \text{ mm Hg} = 760 \text{ torr}$$

$$\text{STP} = 273.15 \text{ K and } 1.0 \text{ atm}$$

$$\text{Ideal gas at STP} = 22.4 \text{ L mol}^{-1}$$

THERMODYNAMICS / ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

q = heat

m = mass

c = specific heat capacity

T = temperature

S° = standard entropy

H° = standard enthalpy

G° = standard Gibbs free energy

n = number of moles

E° = standard reduction potential

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

$$\text{Faraday's constant, } F = 96,485 \text{ coulombs per mole of electrons}$$

$$1 \text{ volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$

CHEMISTRY

Section I

Time—1 hour and 30 minutes

50 Questions

CALCULATORS ARE NOT ALLOWED FOR SECTION I.

Note: For all questions, assume that the temperature is 298 K, the pressure is 1.0 atm, and solutions are aqueous unless otherwise specified.


Directions: Each of the questions or incomplete statements below is followed by four suggested answers or completions. Select the one that is best in each case and then fill in the corresponding circle on the answer sheet.

1. In which of the following processes will ΔS° be negative?

- (A) $\text{C}_2\text{H}_5\text{OH}(l) \rightarrow \text{C}_2\text{H}_5\text{OH}(g)$
- (B) $\text{NaCl}(s) \rightarrow \text{NaCl}(l)$
- (C) $\text{CO}_2(s) \rightarrow \text{CO}_2(g)$
- (D) $\text{Cl}_2(g) \rightarrow \text{Cl}_2(l)$

Solution
0.1 M $\text{HC}_2\text{H}_3\text{O}_2(aq)$
0.1 M $\text{KI}(aq)$
0.1 M $\text{CH}_3\text{OH}(aq)$

2. Of the three solutions listed in the table above, which one, if any, has the greatest electrical conductivity and why?
- (A) 0.1 M $\text{HC}_2\text{H}_3\text{O}_2(aq)$ because its molecules have the most atoms.
 - (B) 0.1 M $\text{KI}(aq)$ because KI completely dissociates in water to produce ions.
 - (C) 0.1 M $\text{CH}_3\text{OH}(aq)$ because its molecules can form hydrogen bonds.
 - (D) All three solutions have the same electrical conductivity because the concentrations are the same.




3. Hydrogen gas reacts with iodine gas at constant temperature in a sealed rigid container. The gases are allowed to reach equilibrium according to the equation above. Which of the following best describes what will happen to the reaction immediately after additional iodine gas is added to the system?

- (A) The rates of both the forward and reverse reactions decrease.
- (B) The rates of both the forward and reverse reactions do not change.
- (C) The rate of the forward reaction becomes greater than the rate of the reverse reaction.
- (D) The rate of the forward reaction becomes less than the rate of the reverse reaction.

	Ionization Energy (kJ/mol)
First	578
Second	1,817
Third	2,745
Fourth	11,577
Fifth	14,842


4. The first five ionization energies of an unknown element are listed in the table above. Which of the following statements correctly identifies the element and cites the evidence supporting the identification?
- (A) Na, because of the large difference between the first and the second ionization energies
 - (B) Al, because of the large difference between the third and fourth ionization energies
 - (C) Si, because the fifth ionization energy has the greatest value
 - (D) P, because a neutral atom of P has five valence electrons



5. The diagram above represents a particle in aqueous solution. Which of the following statements about the particle is correct?
- (A) The particle must be a cation because the negative end of each water molecule is pointed toward it.
 - (B) The particle must be an anion because the positive end of each water molecule is pointed toward it.
 - (C) The charge of the particle cannot be determined because water molecules have no net charge.
 - (D) The charge of the particle cannot be determined because the water molecules are arranged symmetrically and their partial charges cancel.

Compound	Formula	Boiling Point (°C)	Density (g/mL)
Hexane	C_6H_{14}	69	0.66
Octane	C_8H_{18}	126	0.70

6. A student obtains a mixture of the liquids hexane and octane, which are miscible in all proportions. Which of the following techniques would be best for separating the two components of the mixture, and why?
- (A) Filtration, because the different densities of the liquids would allow one to pass through the filter paper while the other would not.
 - (B) Paper chromatography, because the liquids would move along the stationary phase at different rates owing to the difference in polarity of their molecules.
 - (C) Column chromatography, because the higher molar mass of octane would cause it to move down the column faster than hexane.
 - (D) Distillation, because the liquids would boil at different temperatures owing to the difference in strength of their intermolecular forces.







7. The value of the equilibrium constant for the reaction represented above is 1×10^{10} . What is the value of the equilibrium constant for the following reaction?




- (A) 1×10^{-20}
 (B) 1×10^{-10}
 (C) 1×10^{10}
 (D) 1×10^{20}
8. To catalyze a biochemical reaction, an enzyme typically
- (A) drives the reaction to completion by consuming byproducts of the reaction
 (B) binds temporarily to reactant molecules to lower the activation energy of the reaction
 (C) dissociates into additional reactant molecules, thereby increasing the reaction rate
 (D) decomposes and releases energy to increase the number of successful collisions between reactant molecules

9. Of the following diagrams, which best represents the structure of solid KF?



Questions 10-11 refer to the following information.



Standard reduction potentials for the half-reactions associated with the electrochemical cell shown above are given in the table below.

Half-Reaction	E° (V)
$\text{Ag}^{+}(aq) + e^{-} \rightarrow \text{Ag}(s)$	+ 0.80
$\text{Zn}^{2+}(aq) + 2 e^{-} \rightarrow \text{Zn}(s)$	− 0.76

10. Which of the following is the net ionic equation for the overall reaction that occurs as the cell operates?

- (A) $\text{Ag}(s) + \text{Zn}^{2+}(aq) \rightarrow \text{Ag}^{2+}(aq) + \text{Zn}(s)$
- (B) $\text{Ag}^{+}(aq) + \text{Zn}^{2+}(aq) + 3 e^{-} \rightarrow \text{AgZn}(s)$
- (C) $\text{Ag}^{+}(aq) + \text{Zn}(s) \rightarrow \text{Ag}(s) + \text{Zn}^{2+}(aq) + e^{-}$
- (D) $2 \text{Ag}^{+}(aq) + \text{Zn}(s) \rightarrow 2 \text{Ag}(s) + \text{Zn}^{2+}(aq)$

11. What is the value of E° for the cell?

- (A) 0.04 V
- (B) 0.84 V
- (C) 1.56 V
- (D) 2.36 V



12. In a one-person spacecraft, an astronaut exhales 880 g of $\text{CO}_2(g)$ (molar mass 44 g/mol) per day. To prevent the buildup of $\text{CO}_2(g)$ in the spacecraft, a device containing $\text{LiOH}(s)$ is used to remove the $\text{CO}_2(g)$, as represented by the equation above. What mass of $\text{LiOH}(s)$ (molar mass 24 g/mol) is needed to react with all of the $\text{CO}_2(g)$ produced by an astronaut in one day?
- (A) 40. g
(B) 240 g
(C) 480 g
(D) 960 g

Time (s)	[NO ₂] (mol/L)	ln [NO ₂]	$\frac{1}{[\text{NO}_2]}$ (L/mol)
0	0.500	-0.693	2.00
100	0.364	-1.01	2.75
200	0.286	-1.25	3.50
300	0.235	-1.45	4.25



13. The data from a study of the decomposition of NO₂(g) to form NO(g) and O₂(g) are given in the table above. Which of the following rate laws is consistent with the data?

(A) Rate = $k [\text{NO}_2]$

(B) Rate = $k [\text{NO}_2]^2$

(C) Rate = $k \frac{1}{[\text{NO}_2]}$

(D) Rate = $k \frac{1}{[\text{NO}_2]^2}$



14. The photoelectron spectra for H and He are represented above. Which of the following statements best accounts for the fact that the peak on the He spectrum is farther to the left and higher than the peak on the H spectrum?
- (A) He has an additional valence electron in a higher energy level than the valence electron in H.
 - (B) He has a greater nuclear charge than H and an additional electron in the same energy level.
 - (C) He has a completely filled valence shell in which the electrons are a greater distance from the nucleus than the distance between the H nucleus and its electron.
 - (D) It takes longer for the electrons in He to be removed due to the higher nuclear mass of He.

Questions 15-19 are based on the information in the table below.

Compound	Molar Mass	Balanced Equation for the Combustion of One Mole of the Compound	ΔH_{comb}° (kJ/mol _{rxn})
$\begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \\ \text{Ethane} \end{array}$	30. g/mol	$\text{C}_2\text{H}_6(g) + \frac{7}{2}\text{O}_2(g) \rightarrow 2\text{CO}_2(g) + 3\text{H}_2\text{O}(l)$	-1560
$\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\ddot{\text{O}}-\text{H} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \\ \text{Propanol} \end{array}$	60. g/mol	$\text{C}_3\text{H}_8\text{O}(l) + \frac{9}{2}\text{O}_2(g) \rightarrow 3\text{CO}_2(g) + 4\text{H}_2\text{O}(l)$	-2020
Unknown	?	$? + 5\text{O}_2(g) \rightarrow 4\text{CO}_2(g) + 4\text{H}_2\text{O}(l)$	-2240

15. Which of the following is the molecular formula of the unknown compound?

- (A) C_4H_8
- (B) $\text{C}_4\text{H}_8\text{O}$
- (C) $\text{C}_4\text{H}_4\text{O}_2$
- (D) $\text{C}_4\text{H}_8\text{O}_2$

16. Which of the following best explains why the combustion reactions represented in the table are exothermic?

- (A) The number of bonds in the reactant molecules is greater than the number of bonds in the product molecules.
- (B) The number of bonds in the reactant molecules is less than the number of bonds in the product molecules.
- (C) The energy required to break the bonds in the reactants is greater than the energy released in forming the bonds in the products.
- (D) The energy required to break the bonds in the reactants is less than the energy released in forming the bonds in the products.


17. Based on the structural formulas, which of the following identifies the compound that is more soluble in water and best helps to explain why?
- (A) Ethane, because the electron clouds of its molecules are more polarizable than those of propanol.
 - (B) Ethane, because its molecules can fit into the spaces between water molecules more easily than those of propanol can.
 - (C) Propanol, because its molecules have a greater mass than the molecules of ethane have.
 - (D) Propanol, because its molecules can form hydrogen bonds with water molecules but those of ethane cannot.
18. In an experiment, 30.0 g of ethane and 30.0 g of propanol are placed in separate reaction vessels. Each compound undergoes complete combustion with excess $\text{O}_2(\text{g})$. Which of the following best compares the quantity of heat released in each combustion reaction?
- (A) $q_{\text{ethane}} < q_{\text{propanol}}$
 - (B) $q_{\text{ethane}} = q_{\text{propanol}}$
 - (C) $q_{\text{ethane}} > q_{\text{propanol}}$
 - (D) The quantities of heat released in the combustions of ethane and propanol cannot be compared without knowing the specific heat capacity of the compounds.
19. A sealed 10.0 L flask at 400 K contains equimolar amounts of ethane and propanol in gaseous form. Which of the following statements concerning the average molecular speed of ethane and propanol is true?
- (A) The average molecular speed of ethane is less than the average molecular speed of propanol.
 - (B) The average molecular speed of ethane is greater than the average molecular speed of propanol.
 - (C) The average molecular speed of ethane is equal to the average molecular speed of propanol.
 - (D) The average molecular speeds of ethane and propanol cannot be compared without knowing the total pressure of the gas mixture.

20. As a sample of $\text{KNO}_3(s)$ is stirred into water at 25°C , the compound dissolves endothermically. Which of the following best helps to explain why the process is thermodynamically favorable at 25°C ?





- (A) All endothermic processes are thermodynamically favorable.
- (B) Stirring the solution during dissolution adds the energy needed to drive an endothermic process.
- (C) Dissolving the salt decreases the enthalpy of the system.
- (D) Dissolving the salt increases the entropy of the system.

21. A solution is prepared by mixing equal volumes of $0.20\text{ M HC}_2\text{H}_3\text{O}_2$ and $0.40\text{ M NaC}_2\text{H}_3\text{O}_2$. Which of the following correctly describes what occurs if a small amount of $\text{HCl}(aq)$ or $\text{NaOH}(aq)$ is added?

- (A) If $\text{HCl}(aq)$ is added, the pH will increase only slightly because the Cl^- ions will react with $\text{C}_2\text{H}_3\text{O}_2^-$ ions.
- (B) If $\text{HCl}(aq)$ is added, the pH will decrease only slightly because the H^+ ions will react with $\text{C}_2\text{H}_3\text{O}_2^-$ ions.
- (C) If $\text{NaOH}(aq)$ is added, the pH will increase only slightly because the OH^- ions will react with $\text{C}_2\text{H}_3\text{O}_2^-$ ions.
- (D) If $\text{NaOH}(aq)$ is added, the pH will decrease only slightly because the OH^- ions will react with $\text{HC}_2\text{H}_3\text{O}_2$ molecules.




22. A mixture of $\text{H}_2(g)$ and $\text{O}_2(g)$ is placed in a container as represented above. The $\text{H}_2(g)$ and $\text{O}_2(g)$ react to form $\text{H}_2\text{O}(g)$. Which of the following best represents the container after the reaction has gone to completion?




23. After a certain pesticide compound is applied to crops, its decomposition is a first-order reaction with a half-life of 56 days. What is the rate constant, k , for the decomposition reaction?

(A) 0.012 day^{-1}
 (B) 0.018 day^{-1}
 (C) 56 day^{-1}
 (D) 81 day^{-1}




24. Which of the following can be inferred from the diagram above that shows the dependence of potential energy on the internuclear distance between two atoms?

(A) The atoms form a bond with a bond length of 25 pm.
 (B) The atoms form a bond with a bond length of 75 pm.
 (C) The net force between the atoms is attractive at 25 pm.
 (D) The net force between the atoms is attractive at 75 pm.




25. A mixture of two gases, 0.01 mol of $\text{C}_4\text{H}_{10}(\text{g})$ and 0.065 mol of $\text{O}_2(\text{g})$, is pumped into a cylinder with a movable piston, as shown above. The mixture, originally at 200°C and 1.0 atm, is sparked and the reaction represented below occurs.



Which of the following is true after the product gases return to the original temperature and pressure, and why will the change occur? (Assume all gases behave ideally.)

- (A) The piston will be higher than its original position, because the cylinder will contain a greater number of gas molecules.
 (B) The position of the piston will be unchanged, because the total mass of the gases in the cylinder does not change.
 (C) The position of the piston will be unchanged, because the temperature and pressure of the contents of the cylinder remain the same.
 (D) The piston will be lower than its original position, because the product molecules are smaller than the reactant molecules.

Questions 26-28 are based on the following information.



The structure of haloacetic acids, XCH_2COOH (where X is either F, Cl, Br, or I), is shown above. The dissociation constants and molar masses of four haloacetic acids are listed in the table below.

Acid	$\text{p}K_a$	K_a	Molar Mass (g/mol)
Fluoroacetic acid	2.59	2.57×10^{-3}	78.0
Chloroacetic acid	2.87	1.35×10^{-3}	94.5
Bromoacetic acid	2.90	1.26×10^{-3}	138.9
Iodoacetic acid	3.18	6.61×10^{-4}	185.9

26. Which compound, chloroacetic acid or iodoacetic acid, most likely has the lower boiling point, and why?
- (A) Chloroacetic acid, because the London dispersion forces among its molecules are weaker.
- (B) Chloroacetic acid, because the dipole-dipole forces among its molecules are weaker.
- (C) Iodoacetic acid, because the London dispersion forces among its molecules are stronger.
- (D) Iodoacetic acid, because the dipole-dipole forces among its molecules are stronger.
27. An aqueous solution contains small but equal concentrations of both chloroacetic and fluoroacetic acids. Which statement comparing the percent ionizations of the two acids in the solution is true?
- (A) The percent ionization of chloroacetic acid is greater than that of fluoroacetic acid.
- (B) The percent ionization of chloroacetic acid is less than that of fluoroacetic acid.
- (C) The percent ionizations cannot be compared without knowing the concentrations of the two acids.
- (D) The percent ionizations cannot be compared without knowing the pH of the solution.
28. A student titrates 10.0 mL samples of 1.0 M solutions of each of the haloacetic acids with a standard solution of NaOH. Which of the following statements correctly predicts the volume of NaOH(aq) needed to reach the equivalence point?
- (A) Fluoroacetic acid will need the smallest volume of NaOH(aq) to reach the equivalence point.
- (B) Iodoacetic acid will need the smallest volume of NaOH(aq) to reach the equivalence point.
- (C) All of the acids will need the same volume of NaOH(aq) to reach the equivalence point.
- (D) All of the haloacetic acids are weak; therefore none will reach an equivalence point.


29. The value of K_w at 40°C is 3.0×10^{-14} . What is the pH of pure water at 40°C ?
- (A) 3.0
(B) 6.8
(C) 7.0
(D) 7.2
30. A 1.0 mol sample of $\text{He}(g)$ at 25°C is mixed with a 1.0 mol sample of $\text{Xe}(g)$ at 50°C . Which of the following correctly predicts the changes in average kinetic energy and the average speed of the $\text{Xe}(g)$ atoms that will occur as the mixture approaches thermal equilibrium?

	Average Kinetic Energy of Xe Atoms	Average Speed of Xe Atoms
(A)	Will increase	Will increase
(B)	Will increase	Will decrease
(C)	Will decrease	Will increase
(D)	Will decrease	Will decrease

31. Which of the following best helps to explain why $\text{Na}(s)$ is more reactive with water than $\text{Mg}(s)$ is?
- (A) $\text{Na}(s)$ is softer than $\text{Mg}(s)$.
(B) The atomic mass of Na is less than that of Mg.
(C) The Na^+ ion has weaker Coulombic attraction to an anion than the Mg^{2+} ion has.
(D) The first ionization energy of Na is less than that of Mg.
32. At 25°C , enough distilled water is added to a 30.0 mL sample of $\text{HNO}_3(aq)$ with a pH of 4.20 so that the final pH of the diluted solution is 5.20. The volume of distilled water added to the original solution is closest to
- (A) 30.0 mL
(B) 60.0 mL
(C) 270. mL
(D) 300. mL

33. A student observes that the equilibrium constant for a reaction is greater than 1.0 at temperatures below 500 K but less than 1.0 at temperatures above 500 K. What can the student conclude about the values of ΔH° and ΔS° for the reaction? (Assume that ΔH° and ΔS° are independent of temperature.)


- (A) $\Delta H^\circ > 0$ and $\Delta S^\circ > 0$
(B) $\Delta H^\circ > 0$ and $\Delta S^\circ < 0$
(C) $\Delta H^\circ < 0$ and $\Delta S^\circ > 0$
(D) $\Delta H^\circ < 0$ and $\Delta S^\circ < 0$



34. The saturated $\text{CuSO}_4(aq)$ shown above is left uncovered on a lab bench at a constant temperature. As the solution evaporates, 1.0 mL samples of the solution are removed every three days and the $[\text{SO}_4^{2-}]$ in the samples is measured. It is observed that the $[\text{SO}_4^{2-}]$ in the solution did not change over time. Which of the following best helps to explain the observation?
- (A) As the solution evaporates, $\text{Cu}^{2+}(aq)$ and $\text{SO}_4^{2-}(aq)$ leave the beaker along with the water molecules.
(B) As the solution evaporates, the dissolution of $\text{CuSO}_4(s)$ in the beaker decreases.
(C) The evaporation of water is endothermic, so more $\text{CuSO}_4(s)$ dissolves exothermically in the solution, which increases the $[\text{SO}_4^{2-}]$.
(D) As water evaporates, more $\text{CuSO}_4(s)$ precipitates out of the solution in the beaker.

Questions 35-38 refer to the following information.

An experiment is performed to measure the mass percent of $\text{CaCO}_3(s)$ in eggshells. Five different samples of $\text{CaCO}_3(s)$ of known mass react with an excess of $2.0\text{ M HCl}(aq)$ in identical sealed, rigid reaction vessels. The pressure of the gas produced is measured with a pressure gauge attached to the reaction vessel. Since the reaction is exothermic, the reaction system is cooled to its original temperature before the pressure is recorded. The experimental data are used to create the calibration line below.







The experiment is repeated with an eggshell sample, and the experimental data are recorded in the table below.

Mass of eggshell sample	0.200 g
Pressure prior to reaction	0.800 atm
Pressure at completion of reaction	0.870 atm

35. Which of the following equations best represents the species that react and the species that are produced when $\text{CaCO}_3(s)$ and $\text{HCl}(aq)$ are combined?
- (A) $2\text{H}^+(aq) + \text{CaCO}_3(s) \rightarrow \text{Ca}^{2+}(aq) + \text{CO}_3^{2-}(aq) + \text{H}_2(g)$
(B) $2\text{H}^+(aq) + \text{CaCO}_3(s) \rightarrow \text{Ca}^{2+}(aq) + \text{H}_2\text{O}(l) + \text{CO}_2(g)$
(C) $2\text{H}^+(aq) + \text{CaCO}_3(s) \rightarrow \text{Ca}^{2+}(aq) + 2\text{OH}^-(aq) + \text{CO}(g)$
(D) $2\text{HCl}(aq) + \text{CaCO}_3(s) \rightarrow \text{Ca}^{2+}(aq) + \text{H}_2\text{O}(l) + \text{CO}_2(g) + 2\text{Cl}^-(aq)$
36. The mass percent of $\text{CaCO}_3(s)$ in the eggshell sample is closest to
- (A) 30%
(B) 45%
(C) 60%
(D) 75%
37. Another sample of eggshell reacts completely with 4.0 mL of an $\text{HCl}(aq)$ solution of unknown concentration. If the reaction produced 0.095 atm of gas, the concentration of the $\text{HCl}(aq)$ solution was at least
- (A) 0.0020 *M*
(B) 0.050 *M*
(C) 0.50 *M*
(D) 1.0 *M*
38. Which of the following modifications will increase the rate of the reaction the most?
- (A) Using 2.0 *M* $\text{CH}_3\text{COOH}(aq)$ instead of 2.0 *M* $\text{HCl}(aq)$
(B) Cooling the $\text{HCl}(aq)$ to a lower temperature than it was in the original experiment
(C) Reducing the volume of the reaction vessel
(D) Using eggshells that are more finely powdered than those used in the original experiment

39. Samples of $\text{NaF}(s)$ and $\text{NH}_4\text{Cl}(s)$ are dissolved in separate beakers that each contain 100 mL of water. One of the salts produces a slightly acidic solution. Which of the following equations best represents the formation of the slightly acidic solution?






40. Which of the following best explains why the reaction represented above is not observed to occur at room temperature?
- (A) The rate of the reaction is extremely slow because of the relatively small value of ΔG° for the reaction.
 - (B) The entropy of the system decreases because the carbon atoms in graphite are less ordered than those in diamond.
 - (C) The reaction has an extremely large activation energy due to strong three-dimensional bonding among carbon atoms in diamond.
 - (D) The reaction does not occur because it is not thermodynamically favorable.
-



41. The reaction represented above occurs when 2.00×10^{-4} mol of pure acetylsalicylic acid, $\text{C}_9\text{H}_8\text{O}_4$, is completely dissolved in 15.00 mL of water in a flask and titrated to the equivalence point with 0.100 M $\text{NaOH}(\text{aq})$. Which of the following statements about the titration is true at the equivalence point?
- (A) $[\text{C}_9\text{H}_8\text{O}_4]$ is greater than $[\text{C}_9\text{H}_7\text{O}_4^-]$.
 - (B) $[\text{C}_9\text{H}_8\text{O}_4]$ is less than $[\text{C}_9\text{H}_7\text{O}_4^-]$.
 - (C) $[\text{C}_9\text{H}_8\text{O}_4]$ is equal to $[\text{C}_9\text{H}_7\text{O}_4^-]$.
 - (D) The relationship between $[\text{C}_9\text{H}_8\text{O}_4]$ and $[\text{C}_9\text{H}_7\text{O}_4^-]$ cannot be determined without additional information.

Questions 42-44 refer to the following.




The structural formula of the glycinium cation is shown above. Arrows indicate the $\text{p}K_a$ values for the labile protons in the molecule.

42. What is the approximate H–O–C bond angle in the glycinium cation?

(A) 180°
(B) 120°
(C) 105°
(D) 90°


43. Which of the following is true about the geometry of the glycinium cation?

(A) The leftmost C atom and all the atoms directly bonded to it lie in the same plane.
(B) Both C atoms and both O atoms lie in the same plane.
(C) The N–C–C bond angle is 180° .
(D) The geometry around the N atom is planar.



44. Which of the following statements is true about sodium glycinate, represented above?

(A) It is an ionic solid at room temperature.
(B) It is not soluble in water.
(C) It evaporates easily because of the lack of hydrogen bonding.
(D) It dissolves in water to form an acidic solution.




45. Solid carbon tetrachloride, $\text{CCl}_4(s)$, is represented by the diagram above. The attractions between the CCl_4 molecules that hold the molecules together in the solid state are best identified as

(A) polar covalent bonds
 (B) nonpolar covalent bonds
 (C) intermolecular attractions resulting from temporary dipoles
 (D) intermolecular attractions resulting from permanent dipoles


46. Which of the following is a list of the minimum amount of data needed for determining the molar enthalpy of solution of $\text{KCl}(s)$ in pure $\text{H}_2\text{O}(l)$? (Assume that the $\text{KCl}(aq)$ has the same specific heat capacity as pure water and that the initial temperatures of the $\text{KCl}(s)$ and the water are the same.)

(A) Mass of $\text{KCl}(s)$, initial temperature of the water, and final temperature of the solution
 (B) Mass of H_2O , initial temperature of the water, and final temperature of the solution
 (C) Mass of $\text{KCl}(s)$, mass of H_2O , initial temperature of the water, and final temperature of the solution
 (D) Mass of $\text{KCl}(s)$, mass of H_2O , initial temperature of the water, final temperature of the solution, and atmospheric pressure




47. A mixture of $\text{X}_2(g)$, $\text{Y}_2(g)$, and $\text{XY}(g)$ is placed in a previously evacuated, rigid container and allowed to reach equilibrium at a constant temperature, as shown above. Which of the following sets of initial concentrations would lead to the formation of more product as the system moves toward equilibrium?

	$[\text{X}_2]_{\text{initial}}$	$[\text{Y}_2]_{\text{initial}}$	$[\text{XY}]_{\text{initial}}$
(A)	0.40 M	0.40 M	0.20 M
(B)	0.30 M	0.30 M	0.90 M
(C)	0.15 M	0.15 M	0.30 M
(D)	0.10 M	0.10 M	0.20 M



cis-isomer



trans-isomer

48. The structural formulas for two isomers of 1,2-dichloroethene are shown above. Which of the two liquids has the higher equilibrium vapor pressure at 20°C, and why?
- (A) The *cis*-isomer, because it has dipole-dipole interactions, whereas the *trans*-isomer has only London dispersion forces
 - (B) The *cis*-isomer, because it has only London dispersion forces, whereas the *trans*-isomer also has dipole-dipole interactions
 - (C) The *trans*-isomer, because it has dipole-dipole interactions, whereas the *cis*-isomer has only London dispersion forces
 - (D) The *trans*-isomer, because it has only London dispersion forces, whereas the *cis*-isomer also has dipole-dipole interactions



49. The concentration of $\text{F}^{-}(aq)$ in drinking water that is considered to be ideal for promoting dental health is $4.0 \times 10^{-5} M$. Based on the information above, the maximum concentration of $\text{Ca}^{2+}(aq)$ that can be present in drinking water without lowering the concentration of $\text{F}^{-}(aq)$ below the ideal level is closest to
- (A) 0.25 *M*
 - (B) 0.025 *M*
 - (C) $1.6 \times 10^{-6} M$
 - (D) $1.6 \times 10^{-15} M$



50. The formation of $\text{Fe}(s)$ and $\text{O}_2(g)$ from $\text{FeO}(s)$ is not thermodynamically favorable at room temperature. In an effort to make the process favorable, $\text{C}(s)$ is added to the $\text{FeO}(s)$ at elevated temperatures. Based on the information above, which of the following gives the value of K_{eq} and the sign of ΔG° for the reaction represented by the equation below at 1000 K?



	K_{eq}	ΔG°
(A)	1×10^{-38}	Positive
(B)	1×10^{-38}	Negative
(C)	1×10^{26}	Positive
(D)	1×10^{26}	Negative

END OF SECTION I

**IF YOU FINISH BEFORE TIME IS CALLED,
YOU MAY CHECK YOUR WORK ON THIS SECTION.**

DO NOT GO ON TO SECTION II UNTIL YOU ARE TOLD TO DO SO.

MAKE SURE YOU HAVE DONE THE FOLLOWING.

- **PLACED YOUR AP NUMBER LABEL ON YOUR ANSWER SHEET**
- **WRITTEN AND GRIDDED YOUR AP NUMBER CORRECTLY ON YOUR ANSWER SHEET**
- **TAKEN THE AP EXAM LABEL FROM THE FRONT OF THIS BOOKLET AND PLACED IT ON YOUR ANSWER SHEET**

AP[®] Chemistry Exam

SECTION II: Free Response

2019

DO NOT OPEN THIS BOOKLET UNTIL YOU ARE TOLD TO DO SO.

At a Glance

Total Time

1 hour and 45 minutes

Number of Questions

7

Percent of Total Score

50%

Writing Instrument

Either pencil or pen with black or dark blue ink

Electronic Device

Calculator allowed

Suggested Time

Approximately
23 minutes each for
questions 1—3 and
9 minutes each for
questions 4—7

Weight

Approximate weights:
Questions 1—3:
22% each
Questions 4—7:
9% each

IMPORTANT Identification Information

PLEASE PRINT WITH PEN:

1. First two letters of your last name

First letter of your first name

2. Date of birth

Month Day Year

3. Six-digit school code

4. Unless I check the box below, I grant the College Board the unlimited right to use, reproduce, and publish my free-response materials, both written and oral, for educational research and instructional purposes. My name and the name of my school will not be used in any way in connection with my free-response materials. I understand that I am free to mark "No" with no effect on my score or its reporting.

No, I do not grant the College Board these rights. ☐

Instructions

The questions for Section II are printed in this booklet. Pages containing a periodic table and lists containing equations and constants are also printed in this booklet.

You may use the pages that the questions are printed on to organize your answers or for scratch work, but you must write your answers in the areas designated for each response. Only material written in the space provided will be scored.

Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.

Write clearly and legibly. Cross out any errors you make; erased or crossed-out work will not be scored.

Manage your time carefully. You may proceed freely from one question to the next. You may review your responses if you finish before the end of the exam is announced.

Form I
Form Code 4BP4-S

25

DO NOT DETACH FROM BOOK.

PERIODIC TABLE OF THE ELEMENTS																	
1	2																18
1																	2
H 1.008																	He 4.00
3	4															9	10
Li 6.94	Be 9.01															F 19.00	Ne 20.18
11	12															17	18
Na 22.99	Mg 24.30															Cl 35.45	Ar 39.95
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K 39.10	Ca 40.08	Sc 44.96	Ti 47.87	V 50.94	Cr 52.00	Mn 54.94	Fe 55.85	Co 58.93	Ni 58.69	Cu 63.55	Zn 65.38	Ga 69.72	Ge 72.63	As 74.92	Se 78.97	Br 79.90	Kr 83.80
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb 85.47	Sr 87.62	Y 88.91	Zr 91.22	Nb 92.91	Mo 95.95	Tc (97)	Ru 101.1	Rh 102.91	Pd 106.42	Ag 107.87	Cd 112.41	In 114.82	Sn 118.71	Sb 121.76	Te 127.60	I 126.90	Xe 131.29
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs 132.91	Ba 137.33	*La 138.91	Hf 178.49	Ta 180.95	W 183.84	Re 186.21	Os 190.2	Ir 192.2	Pt 195.08	Au 196.97	Hg 200.59	Tl 204.38	Pb 207.2	Bi 208.98	Po (209)	At (210)	Rn (222)
87	88	89	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118
Fr (223)	Ra (226)	†Ac (227)	Rf (267)	Db (270)	Sg (271)	Bh (270)	Hs (277)	Mt (276)	Ds (281)	Rg (282)	Cn (285)	Uut (285)	Fl (289)	Uup (288)	Lv (293)	Uus (294)	Uuo (294)

*Lanthanoid Series													
58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm (145)	62 Sm 150.4	63 Eu 151.97	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.05	71 Lu 174.97
†Actinoid Series													
90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np (237)	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (262)

AP[®] CHEMISTRY EQUATIONS AND CONSTANTS

Throughout the exam the following symbols have the definitions specified unless otherwise noted.

L, mL = liter(s), milliliter(s)
 g = gram(s)
 nm = nanometer(s)
 atm = atmosphere(s)

mm Hg = millimeters of mercury
 J, kJ = joule(s), kilojoule(s)
 V = volt(s)
 mol = mole(s)

ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

E = energy
 ν = frequency
 λ = wavelength

Planck's constant, $h = 6.626 \times 10^{-34}$ J s

Speed of light, $c = 2.998 \times 10^8$ m s⁻¹

Avogadro's number = 6.022×10^{23} mol⁻¹

Electron charge, $e = -1.602 \times 10^{-19}$ coulomb

EQUILIBRIUM

$$K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}, \text{ where } aA + bB \rightleftharpoons cC + dD$$

$$K_p = \frac{(P_C)^c(P_D)^d}{(P_A)^a(P_B)^b}$$

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[H^+], \text{ pOH} = -\log[OH^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[A^-]}{[HA]}$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

Equilibrium Constants

K_c (molar concentrations)

K_p (gas pressures)

K_a (weak acid)

K_b (weak base)

K_w (water)

KINETICS

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

k = rate constant

t = time

$t_{1/2}$ = half-life

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = ^\circ\text{C} + 273$$

$$D = \frac{m}{V}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

Molarity, M = moles of solute per liter of solution

$$A = abc$$

P = pressure

V = volume

T = temperature

n = number of moles

m = mass

M = molar mass

D = density

KE = kinetic energy

v = velocity

A = absorbance

a = molar absorptivity

b = path length

c = concentration

$$\begin{aligned}\text{Gas constant, } R &= 8.314 \text{ J mol}^{-1} \text{ K}^{-1} \\ &= 0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1} \\ &= 62.36 \text{ L torr mol}^{-1} \text{ K}^{-1}\end{aligned}$$

$$1 \text{ atm} = 760 \text{ mm Hg} = 760 \text{ torr}$$

$$\text{STP} = 273.15 \text{ K and } 1.0 \text{ atm}$$

$$\text{Ideal gas at STP} = 22.4 \text{ L mol}^{-1}$$

THERMODYNAMICS / ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

q = heat

m = mass

c = specific heat capacity

T = temperature

S° = standard entropy

H° = standard enthalpy

G° = standard Gibbs free energy

n = number of moles

E° = standard reduction potential

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

$$\text{Faraday's constant, } F = 96,485 \text{ coulombs per mole of electrons}$$

$$1 \text{ volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$

SECTION II BEGINS ON PAGE 6.

CHEMISTRY

Section II

Time—1 hour and 45 minutes

7 Questions

YOU MAY USE YOUR CALCULATOR FOR THIS SECTION.

Directions: Questions 1–3 are long free-response questions that require about 23 minutes each to answer and are worth 10 points each. Questions 4–7 are short free-response questions that require about 9 minutes each to answer and are worth 4 points each.


Write your response in the space provided following each question. Examples and equations may be included in your responses where appropriate. For calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Pay attention to significant figures.




1. Ethanol, $\text{C}_2\text{H}_5\text{OH}$, will combust in air according to the equation above.
 - (a) Is $\text{O}_2(g)$ oxidized in the reaction, or is it reduced? Justify your answer in terms of oxidation numbers.
 - (b) When a sample of $\text{C}_2\text{H}_5\text{OH}$ was combusted, the volume of $\text{CO}_2(g)$ produced was 18.0 L when measured at 21.7°C and 1.03 atm. Determine the number of moles of $\text{CO}_2(g)$ that was produced.
 - (c) Determine the volume of $\text{C}_2\text{H}_5\text{OH}(l)$, in mL, that was combusted to produce the volume of $\text{CO}_2(g)$ collected in part (b). (The density of $\text{C}_2\text{H}_5\text{OH}(l)$ is 0.79 g/mL.)
 - (d) Determine the amount of heat, in kJ, that is released by the combustion reaction.
- The combustion reaction occurred in a closed room containing 5.56×10^4 g of air originally at 21.7°C . Assume that all of the heat produced by the reaction was absorbed by the air (specific heat = $1.005 \text{ J/(g } ^\circ\text{C})$) in the room.
- (e) Determine the final temperature of the air in the room after the combustion.
 - (f) The boiling points of ethanol and dimethyl ether are provided in the following table. Identify the intermolecular force that is most responsible for the difference in boiling point. In terms of this force, explain why the boiling point of ethanol is higher than that of dimethyl ether.

[illegible]


[illegible]



2. Hydrogen peroxide, H_2O_2 , decomposes according to the equation above. This reaction is thermodynamically favorable at room temperature.
- (a) A particulate representation of the reactants is shown below in the box on the left. In the box below on the right, draw the particulate representation of all the molecules that would be produced from these four reactant molecules.




- (b) Shown below is a potential energy diagram for the uncatalyzed decomposition of $\text{H}_2\text{O}_2(aq)$.




- (i) According to the diagram, is the decomposition reaction exothermic or endothermic? Justify your answer.
- (ii) Manganese dioxide, $\text{MnO}_2(s)$, is an insoluble substance that acts as a catalyst for the decomposition reaction. On the diagram above, draw a curve to represent the reaction as it occurs in the presence of $\text{MnO}_2(s)$.

A student investigates the decomposition reaction in the laboratory. The student prepares two small beakers, adding 20.0 mL of 9.77 M $\text{H}_2\text{O}_2(aq)$ to each one. Each beaker is placed on an electronic balance. The student adds 0.10 g of $\text{MnO}_2(s)$ to the second beaker and records the mass of each beaker and its contents at 10-second intervals for one minute. The beakers and the data are shown below.



Beaker 1



Beaker 2


Time (seconds)	Mass of Beaker 1 + $\text{H}_2\text{O}_2(aq)$ (grams)	Mass of Beaker 2 + $\text{H}_2\text{O}_2(aq)$ + $\text{MnO}_2(s)$ (grams)
0	43.09	43.19
10	43.09	43.06
20	43.09	42.94
30	43.09	42.83
40	43.09	42.73
50	43.09	42.65
60	43.09	42.58

- (c) For beaker 2 during the 60-second period, calculate the following.
- The number of moles of $\text{O}_2(g)$ that was produced
 - The mass of $\text{H}_2\text{O}_2(aq)$ that decomposed
- (d) The student continues the experiment for an additional minute. For beaker 2, will the mass of $\text{H}_2\text{O}_2(aq)$ consumed during the second minute be greater than, less than, or equal to the mass of $\text{H}_2\text{O}_2(aq)$ consumed during the first minute? Explain your answer referring to the data in the table.
- (e) Based on the data, the student claims that the catalyzed reaction has zeroth-order kinetics. Do you agree with the student's claim? Justify your answer.
- (f) A second student did the experiment using larger volumes of 9.77 M $\text{H}_2\text{O}_2(aq)$. The student noticed that the reaction in beaker 2 proceeded extremely rapidly, causing some of the liquid to splash out of the beaker onto the lab table. The student claims that as a result of the loss of the liquid from the beaker, the calculated number of moles of $\text{O}_2(g)$ produced is greater than the actual number of moles of $\text{O}_2(g)$ produced during the first 60 seconds. Do you agree with the student? Justify your answer.
- (g) The hydrogen peroxide used in this experiment can be prepared by the reaction of solid ammonium peroxydisulfate, $(\text{NH}_4)_2\text{S}_2\text{O}_8$, with water. The products are hydrogen peroxide (H_2O_2) and ammonium bisulfate (NH_4HSO_4). Write the balanced equation for the reaction.

[illegible]


[illegible]

3. Sulfur trioxide, SO_3 , is commercially produced on a massive scale for use in manufacturing. It is also produced as a by-product of burning fossil fuels and is an air pollutant involved in the formation of acid rain.
- (a) In the box below, complete one of the valid Lewis electron-dot diagrams for the SO_3 molecule. Be sure to include all valence electrons, including lone pairs.




- (b) Predict the shape of the SO_3 molecule based on the diagram you completed above.

Under some conditions, $\text{SO}_3(g)$ decomposes to produce $\text{SO}_2(g)$ and $\text{O}_2(g)$, as represented by the equation below.



- (c) Under what temperature conditions will the reaction be thermodynamically favored: high temperatures, low temperatures, all temperatures, or no temperatures? Justify your answer in terms of the change in enthalpy and the change in entropy during the reaction.

A chemist fills a rigid vessel with $\text{SO}_3(g)$ at a certain temperature until the pressure of $\text{SO}_3(g)$ in the container is 0.83 atm. The $\text{SO}_3(g)$ decomposes as the partial pressures of $\text{SO}_3(g)$, $\text{SO}_2(g)$, and $\text{O}_2(g)$ in the container are monitored over time, as shown in the following graph. A change is made to the system at time t_1 .



(d) Answer the following based on the graph.


- Draw an X on the time axis to show when the system first reached equilibrium.
- Describe the change that was made to the system at time t_1 .
- After the change was made at time t_1 , the partial pressure of $\text{SO}_3(g)$ increased while the partial pressure of $\text{O}_2(g)$ decreased. Explain this observation.

(e) Use the information above to answer the following.


- Write the expression for the equilibrium constant, K_p , for the reaction.
- Determine the value of K_p for the reaction.

[illegible]

[illegible]



4. The reaction between hypochlorous acid and water is represented above.
- (a) Identify one of the conjugate acid-base pairs in the reaction.
- (b) Shown in the graph below is the titration curve that results when 25.0 mL of 0.200 *M* HClO is titrated with 0.200 *M* NaOH. Carefully draw a second curve on the graph that would result from the titration of 25.0 mL of 0.200 *M* HClO₄ with 0.200 *M* NaOH.



- (c) A student proposes creating a buffer by dissolving 0.010 mol of $\text{NaClO}_4(s)$ in 100. mL of 0.100 M HClO_4 . Explain why the resulting solution would not be a buffer.

[illegible]

[illegible]

6. Some building materials contain small capsules filled with paraffin wax to improve the insulating properties of the materials. The melting and freezing of paraffin inside the capsules in the material helps to regulate temperature. Paraffin wax begins to melt at 37°C .
- (a) While the wax is melting, is the net flow of thermal energy from the wax to the surroundings or from the surroundings to the wax? Justify your answer.
- (b) Calculate the amount of thermal energy, in kJ, that is required to melt 15.2 grams of solid paraffin wax when the temperature of the surroundings is above the melting point of paraffin. (The molar mass of paraffin is 282.62 g/mol , and its molar heat of fusion is 48.78 kJ/mol .)
- (c) When paraffin wax melts, its density decreases. Provide a particle-level explanation of this observation.

This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper has a slight shadow on the right side, suggesting it's resting on a surface. There is no handwriting or other markings on the paper.


[illegible]

7. A student is given a solution of phenol red of unknown concentration. Solutions of phenol red are bright pink under basic conditions.

- (a) The student uses a pH probe and determines that the pH of the solution is 11.20. Calculate the hydrogen ion concentration, $[H^+]$, in the solution of phenol red.

The student analyzes a sample of the solution using a spectrophotometer set at a wavelength of 559 nm, the wavelength of maximum absorbance for phenol red. The measured absorbance of the phenol red solution at 559 nm is 0.35.

- (b) Based on the calibration curve shown below, what is the concentration of the solution in micromoles per liter (μM) ?



- (c) If the student mixed 10 mL of distilled water with 10 mL of the sample, would this diluted solution have an absorbance greater than, less than, or equal to the absorbance of the original solution? Justify your answer.
- (d) If the student measured the absorbance of the solution at a wavelength of 650 nm, would the absorbance be greater than, less than, or equal to the absorbance of the solution at 559 nm? Justify your answer.

[illegible]

-25-

STOP

END OF EXAM

**IF YOU FINISH BEFORE TIME IS CALLED,
YOU MAY CHECK YOUR WORK ON THIS SECTION.**

**THE FOLLOWING INSTRUCTIONS APPLY TO THE COVERS OF THE
SECTION II BOOKLET.**

- **MAKE SURE YOU HAVE COMPLETED THE IDENTIFICATION INFORMATION AS REQUESTED ON THE FRONT AND BACK COVERS OF THE SECTION II BOOKLET.**
- **CHECK TO SEE THAT YOUR AP NUMBER LABEL APPEARS IN THE BOX ON THE FRONT COVER.**
- **MAKE SURE YOU HAVE USED THE SAME SET OF AP NUMBER LABELS ON ALL AP EXAMS YOU HAVE TAKEN THIS YEAR.**

Answer Key for AP Chemistry
Practice Exam, Section I

Question 1: D	Question 26: A
Question 2: B	Question 27: B
Question 3: C	Question 28: C
Question 4: B	Question 29: B
Question 5: B	Question 30: D
Question 6: D	Question 31: D
Question 7: A	Question 32: C
Question 8: B	Question 33: D
Question 9: B	Question 34: D
Question 10: D	Question 35: B
Question 11: C	Question 36: D
Question 12: D	Question 37: D
Question 13: B	Question 38: D
Question 14: B	Question 39: C
Question 15: D	Question 40: C
Question 16: D	Question 41: B
Question 17: D	Question 42: C
Question 18: C	Question 43: B
Question 19: B	Question 44: A
Question 20: D	Question 45: C
Question 21: B	Question 46: C
Question 22: B	Question 47: A
Question 23: A	Question 48: D
Question 24: B	Question 49: B
Question 25: A	Question 50: D

Multiple-Choice Section for Chemistry

2019 Course Framework Alignment and Rationales

Question 1

Skill	Learning Objective	Topic
1.A	ENE-4.A	Introduction to Entropy
(A)	Incorrect. ΔS for evaporation is positive because the $\text{C}_2\text{H}_5\text{OH}$ molecules in the gas phase can act independently and are distributed over a greater volume than are the molecules in the liquid phase.	
(B)	Incorrect. ΔS for the melting of $\text{NaCl}(s)$ is positive because ions in fixed positions in the solid phase can now move freely among the other ions in the liquid phase.	
(C)	Incorrect. ΔS for the sublimation of $\text{CO}_2(s)$ is positive because the molecules in the gas phase can act independently and are distributed over a greater volume than are the molecules in the solid phase.	
(D)	Correct. ΔS for the condensation of $\text{Cl}_2(g)$ has a negative value because gas molecules become restricted to a much smaller volume as they condense to the liquid phase.	

Question 2

Skill	Learning Objective	Topic
4.C	SPQ-3.A	Solutions and Mixtures
(A)	Incorrect. Although molecules of $\text{HC}_2\text{H}_3\text{O}_2$ do contain the greatest number of atoms, $\text{HC}_2\text{H}_3\text{O}_2(aq)$ dissociates into H^+ ions and $\text{C}_2\text{H}_3\text{O}_2^-(aq)$ ions to a much-smaller extent than the $\text{K}^+(aq)$ and $\text{I}^-(aq)$ ions in $\text{KI}(aq)$; thus, $\text{HC}_2\text{H}_3\text{O}_2(aq)$ is a weaker conductor than $\text{KI}(aq)$.	
(B)	Correct. $\text{KI}(aq)$ has the greatest conductivity because KI dissociates completely in water, producing mobile $\text{K}^+(aq)$ and $\text{I}^-(aq)$ ions that can move through the solution and carry an electrical current.	
(C)	Incorrect. Although CH_3OH molecules can form hydrogen bonds among themselves and with water molecules, they do not dissociate into charged particles that could carry an electrical current. Thus, $\text{CH}_3\text{OH}(aq)$ is not an electrolyte.	
(D)	Incorrect. Electrical conductivity of solutions depends on whether or not the solutes are electrolytes in addition to the concentrations of solutes.	

Question 3

Skill	Learning Objective	Topic
4.A	TRA-6.B	Direction of Reversible Reactions
(A)	Incorrect. Only the rate of the forward reaction increases.	
(B)	Incorrect. The rate of the forward reaction changes, specifically by increasing.	
(C)	Correct. The introduction of more $I_2(g)$ molecules to the system increases the likelihood of a collision between molecules of $I_2(g)$ and $H_2(g)$, which is a necessary step for the reaction to occur. The greater number of such collisions increases the rate of the forward reaction.	
(D)	Incorrect. The rate of the forward reaction becomes greater than the rate of the reverse reaction.	

Question 4

Skill	Learning Objective	Topic
6.E	SAP-2.A	Periodic Trends
(A)	Incorrect. The difference between the first and second ionization energies is much smaller than the difference between the third and fourth ionization energies, which indicates that the element is Al, not Na.	
(B)	Correct. The increase between the third and fourth ionization energies is much greater than the differences between the other successive ionization energies. This indicates that the fourth electron removed from the atom is in an energy subshell ($2p$) that is substantially lower in energy than the subshell from which the first three electrons were removed ($3s$ and $3p$). This is true for the element Al, which has exactly three electrons in the third ($n = 3$) energy level.	
(C)	Incorrect. Although it is true that the fifth ionization energy has the greatest value, it is the <u>difference</u> between successive ionization energies that provides data for determining the identity of an element. The largest difference occurs between the third and fourth ionization energies, which indicates that the element is Al, not Si.	
(D)	Incorrect. Although it is true that a neutral P atom has five valence electrons, it is the difference between successive ionization energies that provides data for determining the identity of an element. The largest difference occurs between the third and fourth ionization energies, which indicates that the element is Al, not P.	

Question 5

Skill	Learning Objective	Topic
6.C	SAP-5.A	Intermolecular Forces
(A)	Incorrect. The negative ends of the water-molecule dipoles are directed <u>away</u> from the particle, not toward it. The central particle has a negative charge, which identifies it as an anion.	
(B)	Correct. The positive end of the permanent dipole of a water molecule is centered between the two hydrogen atoms. Because all the water molecules are arranged with the positive ends of their dipoles oriented toward the central particle, the central particle must have a negative charge, which identifies it as an anion.	
(C)	Incorrect. Although water molecules have no <u>net</u> charge, the distribution of charge in the water molecule is uneven—water molecules have a permanent dipole. Because all the water molecules are arranged with the positive ends of their dipoles oriented toward the central particle, the central particle must have a negative charge, which identifies it as an anion.	
(D)	Incorrect. The partial charges of the water molecules (which give rise to the dipoles) do not necessarily cancel when the whole system consisting of the central particle and the surrounding water molecules is considered because of random molecular motion. However, the charge of the particle is still easily determined because of the orientation of the water-molecule dipoles.	

Question 6

Skill	Learning Objective	Topic
6.E	SPQ-3.C	Separation of Solutions and Mixtures Chromatography
(A)	Incorrect. Filtration would not work at all because both hexane and octane molecules would pass through the relatively huge pores in the filter paper with the same ease. Also, filtration separates mixtures based on particle size, not density.	
(B)	Incorrect. Paper chromatography separates components of mixtures based on differences between affinities of the components for the paper (stationary phase) and the solvent (mobile phase). Both hexane and octane have nonpolar molecules with similar intermolecular forces that would not lead to sufficiently different affinities to allow separation using this method.	
(C)	Incorrect. The masses of the molecules of the two components of the mixture are not a factor in how fast they would move through a column. Column chromatography separates components of mixtures based on differences between affinities of the components for the stationary phase (e.g., silica gel) and the mobile phase. Both hexane and octane have nonpolar molecules with similar intermolecular forces that would not lead to sufficiently different affinities to allow separation using this method.	
(D)	Correct. The difference in boiling points between the two liquids would permit their separation by the process of distillation. In the vapor above a heated mixture of the two liquids, the compound with the lower boiling point (hexane) would predominate. When that vapor cools as it passes through a condenser, liquid hexane could be collected.	

Question 7

Skill	Learning Objective	Topic
5.F	TRA-7.D	Properties of the Equilibrium Constant
(A)	Correct. The second equation can be obtained by simultaneously reversing the direction of the first equation and multiplying each of the coefficients in the equation by 2. The effect of the first transformation is to invert the value of K and the effect of the second transformation is to raise K to the power of 2. Therefore, the value of K for the second equation should have the value $\left(\frac{1}{K}\right)^2 = \frac{1}{K^2}$. Thus, the value of K for the second equation is $\left(\frac{1}{1 \times 10^{10}}\right)^2 = \frac{1}{1 \times 10^{20}} = 1 \times 10^{-20}$.	
(B)	Incorrect. This choice is consistent with failing to recognize that the coefficients of the equation were multiplied by 2.	
(C)	Incorrect. This choice makes the incorrect assumption that the value of K stays constant.	
(D)	Incorrect. This choice is consistent with failing to recognize that the equation was reversed.	

Question 8

Skill	Learning Objective	Topic
4.C	ENE-1.A	Catalysis
(A)	Incorrect. An enzyme does not function by consuming byproducts of the reaction. An enzyme is a catalyst, and as such it provides a different, lower energy pathway for the reaction to occur. The enzyme temporarily binds to the reactant or reactants, allowing the reaction to occur with less input of energy; then the enzyme is regenerated by the release of the product or products.	
(B)	Correct. An enzyme is a catalyst, and as such it provides a different, lower energy pathway for the reaction to occur. The enzyme temporarily binds to the reactant or reactants, allowing the reaction to occur with less input of energy; then the enzyme is regenerated by the release of the product or products.	
(C)	Incorrect. An enzyme does not function by breaking up into more reactant particles. An enzyme is a catalyst, and as such it provides a different, lower energy pathway for the reaction to occur. The enzyme temporarily binds to the reactant or reactants, allowing the reaction to occur with less input of energy; then the enzyme is regenerated by the release of the product or products.	
(D)	Incorrect. An enzyme does not function by decomposing to produce energy. Enzymes are not used up as a reaction occurs, instead they are constantly regenerated. An enzyme is a catalyst, and as such it provides a different, lower energy pathway for the reaction to occur. The enzyme temporarily binds to the reactant or reactants, allowing the reaction to occur with less input of energy; then the enzyme is regenerated by the release of the product or products.	

Question 9

Skill	Learning Objective	Topic
1.A	SAP-3.C	Structure of Ionic Solids
(A)	Incorrect. This diagram shows layers of alternating pairs of ions of the same charge. The repulsions between ions of the same charge in this arrangement would be very large and would not represent a stable configuration. An arrangement in which each ion of one charge was surrounded by ions of the opposite charge would be the most energetically favorable.	
(B)	Correct. This diagram shows ions as being tightly packed surrounded by ions of the opposite charge, which is true in a crystal of an ionic compound such as KF. This arrangement represents the lowest energy state attainable by the constituent ions.	
(C)	Incorrect. This diagram shows ions of the same charge in alternating layers. The repulsions among the ions in the same layer in this arrangement would be very large and would not represent a stable configuration. An arrangement in which each ion of one charge was surrounded by ions of the opposite charge would be the most energetically favorable.	
(D)	Incorrect. This diagram shows ions widely dispersed and in a random arrangement, which could better represent KF in the gas phase, not solid KF.	

Question 10

Skill	Learning Objective	Topic
5.E	TRA-1.B	Net Ionic Equations
(A)	Incorrect. The species $\text{Ag}^{2+}(\text{aq})$ does not exist and is not present in the information provided.	
(B)	Incorrect. This equation incorrectly shows $\text{Ag}^+(\text{aq})$ and $\text{Zn}^{2+}(\text{aq})$ as reactants. In the cell reaction, $\text{Zn}^{2+}(\text{aq})$ is a product, not a reactant.	
(C)	Incorrect. This equation is missing one $\text{Ag}^+(\text{aq})$ on the left side and one $\text{Ag}(\text{s})$ on the right side, and there is an extraneous electron on the right side.	
(D)	Correct. A thermodynamically favorable reaction occurs when aqueous silver ion, $\text{Ag}^+(\text{aq})$, oxidizes zinc metal, $\text{Zn}(\text{s})$, as represented by the equation $2 \text{Ag}^+(\text{aq}) + \text{Zn}(\text{s}) \rightarrow 2 \text{Ag}(\text{s}) + \text{Zn}^{2+}(\text{aq}).$	

Question 11

Skill	Learning Objective	Topic
5.F	ENE-6.B	Cell Potential and Free Energy
(A)	Incorrect. This choice represents the simple addition of the E° value for the two half-reactions.	
(B)	Incorrect. This choice represents the result of adding the E° value for the first half-reaction to the sum of the E° values of the two half-reactions.	
(C)	Correct. The correct reaction is obtained by doubling the first half-reaction in the table (but not doubling the value of E°) and adding the reverse of the second half-reaction (while reversing the sign of its E°). Thus, $E^\circ_{\text{cell}} = 0.80 - (-0.76) = 1.56 \text{ V}$.	
(D)	Incorrect. This choice represents the result of improperly doubling the E° value of the first half-reaction before adding it to the E° value of the second half-reaction.	

Question 12

Skill	Learning Objective	Topic
5.F	SPQ-4.A	Stoichiometry
(A)	Incorrect. In one day the astronaut exhales $880\text{ g} = 20\text{ mol CO}_2(g)$. According to the balanced equation, the amount of $\text{LiOH}(s)$ needed to react with $20\text{ mol CO}_2(g)$ would be $40\text{ mol LiOH}(s)$. The mass of $40\text{ mol LiOH}(s)$ is $40\text{ mol} \times 24\text{ g/mol} = 960\text{ g LiOH}(s)$.	
(B)	Incorrect. In one day the astronaut exhales $880\text{ g} = 20\text{ mol CO}_2(g)$. According to the balanced equation, the amount of $\text{LiOH}(s)$ needed to react with $20\text{ mol CO}_2(g)$ would be $40\text{ mol LiOH}(s)$. The mass of $40\text{ mol LiOH}(s)$ is $40\text{ mol} \times 24\text{ g/mol} = 960\text{ g LiOH}(s)$.	
(C)	Incorrect. In one day the astronaut exhales $880\text{ g} = 20\text{ mol CO}_2(g)$. According to the balanced equation, the amount of $\text{LiOH}(s)$ needed to react with $20\text{ mol CO}_2(g)$ would be $40\text{ mol LiOH}(s)$. The mass of $40\text{ mol LiOH}(s)$ is $40\text{ mol} \times 24\text{ g/mol} = 960\text{ g LiOH}(s)$.	
(D)	Correct. In one day the astronaut exhales $880\text{ g} = 20\text{ mol CO}_2(g)$. According to the balanced equation, the amount of $\text{LiOH}(s)$ needed to react with $20\text{ mol CO}_2(g)$ would be $40\text{ mol LiOH}(s)$. The mass of $40\text{ mol LiOH}(s)$ is $40\text{ mol} \times 24\text{ g/mol} = 960\text{ g LiOH}(s)$.	

Question 13

Skill	Learning Objective	Topic
5.B	TRA-3.C	Concentration Changes Over Time
(A)	Incorrect. The data show that the quantity $\frac{1}{[\text{NO}_2]}$ increases by the same amount (0.75L/mol) every 100 seconds, which means that plotting $\frac{1}{[\text{NO}_2]}$ versus time yields a straight line. According to the integrated rate laws, this result is consistent with the reaction being second order. Therefore, the rate law takes the form $rate = k[\text{NO}_2]^2$.	
(B)	Correct. The data show that the quantity $\frac{1}{[\text{NO}_2]}$ increases by the same amount (0.75L/mol) every 100 seconds, which means that plotting $\frac{1}{[\text{NO}_2]}$ versus time yields a straight line. According to the integrated rate laws, this result is consistent with the reaction being second order. Therefore, the rate law takes the form $rate = k[\text{NO}_2]^2$.	
(C)	Incorrect. The data show that the quantity $\frac{1}{[\text{NO}_2]}$ increases by the same amount (0.75L/mol) every 100 seconds, which means that plotting $\frac{1}{[\text{NO}_2]}$ versus time yields a straight line. According to the integrated rate laws, this result is consistent with the reaction being second order. Therefore, the rate law takes the form $rate = k[\text{NO}_2]^2$.	
(D)	Incorrect. The data show that the quantity $\frac{1}{[\text{NO}_2]}$ increases by the same amount (0.75L/mol) every 100 seconds, which means that plotting $\frac{1}{[\text{NO}_2]}$ versus time yields a straight line. According to the integrated rate laws, this result is consistent with the reaction being second order. Therefore, the rate law takes the form $rate = k[\text{NO}_2]^2$.	

Question 14

Skill	Learning Objective	Topic
6.C	SAP-1.B	Photoelectron Spectroscopy
(A)	Incorrect. The two valence electrons in He are in the <u>same</u> (1s) energy level as that occupied by the valence electron of H. However, the He nucleus has two protons. The greater nuclear charge of He increases the attraction of its two electrons to the nucleus, increasing their binding energy, which moves the peak to the left. The peak is twice the height of the peak for H because there are twice as many electrons in the 1s orbital of the He atom than in the 1s orbital of the H atom.	
(B)	Correct. The He nucleus has two protons. The greater nuclear charge of He increases the attraction of its two electrons to the nucleus, increasing their binding energy, which moves the peak to the left. The peak is twice the height of the peak for H because there are twice as many electrons in the 1s orbital of the He atom than in the 1s orbital of the H atom.	
(C)	Incorrect. The distance of the valence electrons from the nucleus is less in He than in H, not greater. However, the He nucleus has two protons. The greater nuclear charge of He increases the attraction of its two electrons to the nucleus, increasing their binding energy, which moves the peak to the left. The peak is twice the height of the peak for H because there are twice as many electrons in the 1s orbital of the He atom than in the 1s orbital of the H atom.	
(D)	Incorrect. The masses of the two atoms have nothing to do with the amount of time it takes to remove an electron from either one. However, the He nucleus has two protons. The greater nuclear charge of He increases the attraction of its two electrons to the nucleus, increasing their binding energy, which moves the peak to the left. The peak is twice the height of the peak for H because there are twice as many electrons in the 1s orbital of the He atom than in the 1s orbital of the H atom.	

Question 15

Skill	Learning Objective	Topic
5.E	SPQ-4.A	Stoichiometry
(A)	Incorrect. By the law of conservation of mass, the missing atoms on the left side of the equation must be 4 carbon atoms, 8 hydrogen atoms, and 2 oxygen atoms. Therefore, the molecular formula of the unknown compound is $C_4H_8O_2$.	
(B)	Incorrect. By the law of conservation of mass, the missing atoms on the left side of the equation must be 4 carbon atoms, 8 hydrogen atoms, and 2 oxygen atoms. Therefore, the molecular formula of the unknown compound is $C_4H_8O_2$.	
(C)	Incorrect. By the law of conservation of mass, the missing atoms on the left side of the equation must be 4 carbon atoms, 8 hydrogen atoms, and 2 oxygen atoms. Therefore, the molecular formula of the unknown compound is $C_4H_8O_2$.	
(D)	Correct. By the law of conservation of mass, the missing atoms on the left side of the equation must be 4 carbon atoms, 8 hydrogen atoms, and 2 oxygen atoms. Therefore, the molecular formula of the unknown compound is $C_4H_8O_2$.	

Question 16

Skill	Learning Objective	Topic
6.E	ENE-3.A	Bond Enthalpies
(A)	Incorrect. The actual number of bonds in products and reactants, per se, does not control the exothermicity or endothermicity of the reaction. It is the sum of the <u>energies</u> of the bonds that matters. In an exothermic reaction, more energy is released as new bonds in the product molecules form than is required to break the bonds in the reactant molecules.	
(B)	Incorrect. The actual number of bonds in products and reactants, per se, does not control the exothermicity or endothermicity of the reaction. It is the sum of the <u>energies</u> of the bonds that matters. In an exothermic reaction, more energy is released as new bonds in the product molecules form than is required to break the bonds in the reactant molecules.	
(C)	Incorrect. If this were true, the reaction would be endothermic, not exothermic.	
(D)	Correct. In an exothermic reaction, more energy is released as new bonds in the product molecules form than is required to break the bonds in the reactant molecules.	

Question 17

Skill	Learning Objective	Topic
6.E	SAP-5.A	Intermolecular Forces
(A)	Incorrect. The ethane molecule is smaller (14 valence electrons) than the propanol molecule (26 valence electrons); thus, ethane has a smaller, <u>less</u> polarizable electron cloud than propanol has. Also, polarizability of electron clouds is more of a factor in determining London dispersion forces, which are generally weaker than dipole-dipole forces such as hydrogen bonding, which makes propanol more soluble in water than ethane is.	
(B)	Incorrect. Although ethane molecules are smaller than propanol molecules, solubility is affected much more by intermolecular forces between solute and solvent molecules. Propanol molecules, which contain a hydroxyl (-OH) group, can interact more strongly with the polar solvent (H_2O) molecules than ethane molecules, which are nonpolar, can interact with the polar solvent (H_2O) molecules. Thus, the solubility of propanol in water is greater than the solubility of ethane in water.	
(C)	Incorrect. Although propanol molecules have more mass than ethane molecules have, mass itself is not a factor in solubility, which is affected much more by intermolecular forces between solute and solvent molecules. Propanol molecules, which contain a hydroxyl (-OH) group, can interact more strongly with the polar solvent (H_2O) molecules than ethane molecules, which are nonpolar, can interact with the polar solvent (H_2O) molecules. Thus, the solubility of propanol in water is greater than the solubility of ethane in water.	
(D)	Correct. Solubility depends on the nature and strengths of the intermolecular forces between solute and solvent molecules. Propanol molecules, which contain a hydroxyl (-OH) group, can interact more strongly with the polar solvent (H_2O) molecules than ethane molecules, which are nonpolar, can interact with the polar solvent (H_2O) molecules. Thus, the solubility of propanol in water is greater than the solubility of ethane in water.	

Question 18

Skill	Learning Objective	Topic
5.F	ENE-2.F	Introduction to Enthalpy of Reaction
(A)	Incorrect. Although the molar heat of combustion of propanol is greater than that of ethane, 30.0 g of ethane represents one mole and 30.0 g of propanol represents one-half of a mole. According to the data in the table, the combustion of the ethane releases 1560 kJ, whereas the combustion of the propanol releases only one-half of 2020 kJ, or 1010 kJ. Because $1560 \text{ kJ} > 1010 \text{ kJ}$, $q_{\text{ethane}} > q_{\text{propanol}}$.	
(B)	Incorrect. Although the molar heat of combustion of propanol is greater than that of ethane, 30.0 g of ethane represents one mole and 30.0 g of propanol represents one-half of a mole. According to the data in the table, the combustion of the ethane releases 1560 kJ, whereas the combustion of the propanol releases only one-half of 2020 kJ, or 1010 kJ. Because $1560 \text{ kJ} > 1010 \text{ kJ}$, $q_{\text{ethane}} > q_{\text{propanol}}$.	
(C)	Correct. Although the molar heat of combustion of propanol is greater than that of ethane, 30.0 g of ethane represents one mole and 30.0 g of propanol represents one-half of a mole. According to the data in the table, the combustion of the ethane releases 1560 kJ, whereas the combustion of the propanol releases only one-half of 2020 kJ, or 1010 kJ. Because $1560 \text{ kJ} > 1010 \text{ kJ}$, $q_{\text{ethane}} > q_{\text{propanol}}$.	
(D)	Incorrect. The specific heat capacities of the compounds are not needed to calculate the heat released by their combustion. Although the molar heat of combustion of propanol is greater than that of ethane, 30.0 g of ethane represents one mole and 30.0 g of propanol represents one-half of a mole. According to the data in the table, the combustion of the ethane releases 1560 kJ, whereas the combustion of the propanol releases only one-half of 2020 kJ, or 1010 kJ. Because $1560 \text{ kJ} > 1010 \text{ kJ}$, $q_{\text{ethane}} > q_{\text{propanol}}$.	

Question 19

Skill	Learning Objective	Topic
4.A	SAP-7.B	Kinetic Molecular Theory
(A)	Incorrect. Since both gases are at the same temperature (400 K), the average kinetic energies of the ethane and propanol molecules are equal. However, because ethane molecules are less massive than propanol molecules, in order to have the same average kinetic energy $\left(\frac{1}{2}mv^2\right)$, the average molecular speed, v , of the ethane molecules must be <u>greater</u> than (not less than) that of the propanol molecules.	
(B)	Correct. Since both gases are at the same temperature (400 K), the average kinetic energies of the ethane and propanol molecules are equal. However, because ethane molecules are less massive than propanol molecules, in order to have the same average kinetic energy $\left(\frac{1}{2}mv^2\right)$, the average molecular speed, v , of the ethane molecules must be greater than that of the propanol molecules.	
(C)	Incorrect. Since both gases are at the same temperature (400 K), the average kinetic energies of the gases are equal. However, because ethane molecules are less massive than propanol molecules, in order to have the same average kinetic energy $\left(\frac{1}{2}mv^2\right)$, the average molecular speed, v , of the ethane molecules must be greater than (not the same as) that of the propanol molecules.	
(D)	Incorrect. Pressure is irrelevant to the issue of comparing average molecular speeds, and it is possible to compare the average molecular speeds of the two gases. Since both gases are at the same temperature (400 K), the average kinetic energies of the gases are equal. However, because ethane molecules are less massive than propanol molecules, in order to have the same average kinetic energy $\left(\frac{1}{2}mv^2\right)$, the average molecular speed, v , of the ethane molecules must be greater than that of the propanol molecules.	

Question 20

Skill	Learning Objective	Topic
6.D	ENE-4.C	Gibbs Free-Energy and Thermodynamic Favorability
(A)	Incorrect. In fact, most endothermic processes are thermodynamically unfavorable. Because the dissolution process is endothermic, ΔH for the process is positive. The fact that the process occurs indicates that the process is favorable at 25°C; thus, ΔG for the process is negative. Algebraically, since $\Delta G = \Delta H - T\Delta S$, $\Delta G < 0$ implies that $\Delta S > 0$.	
(B)	Incorrect. Stirring can supply a small amount of energy, but the process is driven by an increase in entropy, not enabled by supplying energy for the process to occur. Because the dissolution process is endothermic, ΔH for the process is positive. The fact that the process occurs indicates that the process is favorable at 25°C; thus, ΔG for the process is negative. Algebraically, since $\Delta G = \Delta H - T\Delta S$, $\Delta G < 0$ implies that $\Delta S > 0$.	
(C)	Incorrect. This contradicts the given statement that the salt dissolves endothermically. Because the dissolution process is endothermic, ΔH for the process is positive. The fact that the process occurs indicates that the process is favorable at 25°C; thus, ΔG for the process is negative. Algebraically, since $\Delta G = \Delta H - T\Delta S$, $\Delta G < 0$ implies that $\Delta S > 0$.	
(D)	Correct. Because the dissolution process is endothermic, ΔH for the process is positive. The fact that the process occurs indicates that the process is favorable at 25°C; thus, ΔG for the process is negative. Algebraically, since $\Delta G = \Delta H - T\Delta S$, $\Delta G < 0$ implies that $\Delta S > 0$.	

Question 21

Skill	Learning Objective	Topic
6.E	SAP-10.B	Properties of Buffers
(A)	Incorrect. The anions Cl^- and $\text{C}_2\text{H}_3\text{O}_2^-$ do not react with each other. Furthermore, adding H^+ to a solution would lower its pH, not increase it.	
(B)	Correct. The $\text{C}_2\text{H}_3\text{O}_2^-$ ion, as the anion of a weak acid, is itself a fairly strong base. So the addition of a small amount of H^+ ions will result in most of them reacting with $\text{C}_2\text{H}_3\text{O}_2^-$ ions to form the neutral species $\text{HC}_2\text{H}_3\text{O}_2$. In this way, the pH change in the solution is mitigated, and the pH decreases only slightly.	
(C)	Incorrect. The increase in pH due to the addition of a small amount of OH^- ions will be mitigated because most of the added OH^- anions will react with $\text{HC}_2\text{H}_3\text{O}_2$ molecules, <u>not</u> $\text{C}_2\text{H}_3\text{O}_2^-$ ions, present in the solution.	
(D)	Incorrect. The pH of the solution will <u>increase</u> , not decrease, only slightly because most of the added OH^- anions will react with $\text{HC}_2\text{H}_3\text{O}_2$ molecules present in the solution.	

Question 22

Skill	Learning Objective	Topic
5.E	TRA-1.C	Representations of Reactions
(A)	Incorrect. This diagram violates the law of conservation of matter: four O atoms are missing. The reactants are initially present in equal amounts: four molecules each of H_2 and O_2. In the reaction product, H_2O, there are twice as many H atoms as O atoms; therefore, H_2 is the limiting reactant. When the reaction progresses to completion, all the H_2 molecules will be consumed, four H_2O molecules will be produced, and two O_2 molecules will remain unreacted.	
(B)	Correct. The reactants are initially present in equal amounts: four molecules each of H_2 and O_2. In the reaction product, H_2O, there are twice as many H atoms as O atoms; therefore, H_2 is the limiting reactant. When the reaction progresses to completion, all the H_2 molecules will be consumed, four H_2O molecules will be produced, and two O_2 molecules will remain unreacted, as shown in the diagram for this option.	
(C)	Incorrect. Although the stoichiometry is correct in this diagram, oxygen is represented by four single O atoms instead of two O_2 molecules. If the conditions corresponded to the presence of dissociated O_2 molecules, then the water molecules would also be dissociated.	
(D)	Incorrect. This diagram violates the law of conservation of matter: there are eight extra H atoms present in the products. The reactants are initially present in equal amounts: four molecules each of H_2 and O_2. In the reaction product, H_2O, there are twice as many H atoms as O atoms; therefore, H_2 is the limiting reactant. When the reaction progresses to completion, all the H_2 molecules will be consumed, four H_2O molecules will be produced, and two O_2 molecules will remain unreacted.	

Question 23

Skill	Learning Objective	Topic
5.F	TRA-3.C	Concentration Changes Over Time
(A)	Correct. For a first-order reaction, the half-life $\left(t_{\frac{1}{2}}\right)$ is related to the rate constant (k) by the equation $t_{\frac{1}{2}} = \frac{0.693}{k}$. Rearranging the equation, $k = \frac{0.693}{t_{\frac{1}{2}}} = \frac{0.693}{56 \text{ days}} = 0.012 \text{ day}^{-1}$.	
(B)	Incorrect. For a first-order reaction, the half-life $\left(t_{\frac{1}{2}}\right)$ is related to the rate constant (k) by the equation $t_{\frac{1}{2}} = \frac{0.693}{k}$. Rearranging the equation, $k = \frac{0.693}{t_{\frac{1}{2}}} = \frac{0.693}{56 \text{ days}} = 0.012 \text{ day}^{-1}$.	
(C)	Incorrect. For a first-order reaction, the half-life $\left(t_{\frac{1}{2}}\right)$ is related to the rate constant (k) by the equation $t_{\frac{1}{2}} = \frac{0.693}{k}$. Rearranging the equation, $k = \frac{0.693}{t_{\frac{1}{2}}} = \frac{0.693}{56 \text{ days}} = 0.012 \text{ day}^{-1}$.	
(D)	Incorrect. For a first-order reaction, the half-life $\left(t_{\frac{1}{2}}\right)$ is related to the rate constant (k) by the equation $t_{\frac{1}{2}} = \frac{0.693}{k}$. Rearranging the equation, $k = \frac{0.693}{t_{\frac{1}{2}}} = \frac{0.693}{56 \text{ days}} = 0.012 \text{ day}^{-1}$.	

Question 24

Skill	Learning Objective	Topic
5.D	SAP-3.B	Intramolecular Force and Potential Energy
(A)	Incorrect. At 25 pm the net force between the atoms is repulsive, and a bond is not formed. Rather, the minimum potential energy of the system corresponds to an internuclear distance of 75 pm, at which the bond is stable.	
(B)	Correct. The minimum potential energy occurs at an internuclear distance of 75 pm, which corresponds to the length of the stable bond that forms between the two atoms. If the atoms were any closer to each other, the net force would be repulsive. Likewise, if the atoms were farther from each other, the net force would be attractive. At 75 pm, the net force between the two atoms is zero.	
(C)	Incorrect. At an internuclear distance of 25 pm, the net force between the atoms is repulsive. A stable bond corresponds to an internuclear distance of 75 pm, where the potential energy of the system is as low as possible and the net force between the atoms is zero.	
(D)	Incorrect. The net force between the atoms at an internuclear distance of 75 pm is zero. At this distance, the attractive forces between the atoms are exactly balanced by the repulsive forces.	

Question 25

Skill	Learning Objective	Topic
5.C	SAP-7.A	Ideal Gas Law
(A)	Correct. As the balanced equation indicates, more moles of gas (18) are produced than the number of moles of gas consumed (15) as the reaction proceeds. Therefore, when the reaction is complete and the temperature and pressure are returned to their original values, the volume of gases in the cylinder must be greater (V is proportional to n when both T and P are held constant). Thus, the position of the piston in the cylinder will be higher than its original position.	
(B)	Incorrect. Although mass is conserved during the reaction, as the balanced equation indicates, more moles of gas (18) are produced than the number of moles of gas consumed (15) as the reaction proceeds. Therefore, when the reaction is complete and the temperature and pressure are returned to their original values, the volume of gases in the cylinder must be greater (V is proportional to n when both T and P are held constant). Thus, the position of the piston in the cylinder will be higher than its original position.	
(C)	Incorrect. Although it is true that the temperature and pressure of the gases in the cylinder before and after the reaction are the same, as the balanced equation indicates, more moles of gas (18) are produced than the number of moles of gas consumed (15) as the reaction proceeds. Therefore, when the reaction is complete and the temperature and pressure are returned to their original values, the volume of gases in the cylinder must be greater (V is proportional to n when both T and P are held constant). Thus, the position of the piston in the cylinder will be higher than its original position.	
(D)	Incorrect. The relative sizes of the reactant and product molecules are not relevant because gas behavior is modeled on the gas particles having negligible size compared to the size of interparticle distances within the gas. What is relevant is that, as the balanced equation indicates, more moles of gas (18) are produced than the number of moles of gas consumed (15) as the reaction proceeds. Therefore, when the reaction is complete and the temperature and pressure are returned to their original values, the volume of gases in the cylinder must be greater (V is proportional to n when both T and P are held constant). Thus, the position of the piston in the cylinder will be higher than its original position.	

Question 26

Skill	Learning Objective	Topic
6.E	SAP-5.A	Intermolecular Forces
(A)	Correct. The chlorine atom has fewer occupied shells than the iodine atom. The chloroacetic acid molecule thus has a smaller, less polarizable electron cloud than the iodoacetic acid molecule, so chloroacetic acid has weaker London dispersion forces than iodoacetic acid. The weaker the intermolecular forces, the lower the boiling point.	
(B)	Incorrect. The chloroacetic acid molecule and the iodoacetic acid molecule have the same structure and similar dipole moments. The chloroacetic acid molecule has a smaller, less polarizable electron cloud than the iodoacetic acid molecule, so chloroacetic acid has weaker London dispersion forces and a lower boiling point than iodoacetic acid.	
(C)	Incorrect. Chloroacetic acid has the lower boiling point. The chloroacetic acid molecule has a smaller, less polarizable electron cloud than the iodoacetic acid molecule, so chloroacetic acid has weaker London dispersion forces than iodoacetic acid. The weaker the intermolecular forces, the lower the boiling point.	
(D)	Incorrect. Chloroacetic acid has the lower boiling point. Chloroacetic acid has weaker London dispersion forces than iodoacetic acid because the chloroacetic acid molecule has a smaller, less polarizable electron cloud than the iodoacetic acid molecule. The weaker the intermolecular forces, the lower the boiling point.	

Question 27

Skill	Learning Objective	Topic
4.A	SAP-9.C	Weak Acid and Base Equilibria
(A)	Incorrect. When the acids are at equal concentrations, the percent ionization of chloroacetic acid is <u>less</u> than that of fluoroacetic acid. Fluoroacetic acid has a larger K_a than chloroacetic acid does, so it is a stronger acid and has a larger percent ionization in aqueous solution. The equilibrium-constant expressions for the acids still apply when the two acids are combined, so the percent ionization of fluoroacetic acid will still be greater than that of chloroacetic acid in a mixture of equal concentrations of the acids.	
(B)	Correct. Fluoroacetic acid has a larger K_a than chloroacetic acid does, so it is a stronger acid and at equal concentrations has a larger percent ionization in aqueous solution. The equilibrium-constant expressions for the acids still apply when the two acids are combined, so the percent ionization of fluoroacetic acid will still be greater than that of chloroacetic acid in a mixture of equal concentrations of the acids.	
(C)	Incorrect. The concentrations of the acids are equal, and the concentration is not needed to compare percent ionization. Fluoroacetic acid has a larger K_a than chloroacetic acid does, so it is a stronger acid and at equal concentrations has a larger percent ionization in aqueous solution. The equilibrium-constant expressions for the acids still apply when the two acids are combined, so the percent ionization of fluoroacetic acid will still be greater than that of chloroacetic acid in a mixture of equal concentrations of the acids.	
(D)	Incorrect. The pH is not needed to compare percent ionization. Fluoroacetic acid has a larger K_a than chloroacetic acid does, so it is a stronger acid and at equal concentrations has a larger percent ionization in aqueous solution. The equilibrium-constant expressions for the acids still apply when the two acids are combined, so the percent ionization of fluoroacetic acid will still be greater than that of chloroacetic acid in a mixture of equal concentrations of the acids.	

Question 28

Skill	Learning Objective	Topic
4.A	SPQ-4.B	Introduction to Titration
(A)	Incorrect. At the equivalence point, the number of moles of each monoprotic acid used is equal to the number of moles of $\text{NaOH}(aq)$ added. Since equimolar amounts of the acids are used and the same $\text{NaOH}(aq)$ solution is used for all four titrations, the volume of $\text{NaOH}(aq)$ needed to reach the equivalence point is the same for all four titrations.	
(B)	Incorrect. At the equivalence point, the number of moles of each monoprotic acid used is equal to the number of moles of $\text{NaOH}(aq)$ added. Since equimolar amounts of the acids are used and the same $\text{NaOH}(aq)$ solution is used for all four titrations, the volume of $\text{NaOH}(aq)$ needed to reach the equivalence point is the same for all four titrations.	
(C)	Correct. At the equivalence point, the number of moles of each monoprotic acid used is equal to the number of moles of $\text{NaOH}(aq)$ added. Since equimolar amounts of the acids are used and the same $\text{NaOH}(aq)$ solution is used for all four titrations, the volume of $\text{NaOH}(aq)$ needed to reach the equivalence point is the same for all four titrations.	
(D)	Incorrect. When a weak acid is titrated with a strong base, the titration does have an equivalence point. At the equivalence point, the number of moles of each monoprotic acid used is equal to the number of moles of $\text{NaOH}(aq)$ added. Since equimolar amounts of the acids are used and the same $\text{NaOH}(aq)$ solution is used for all four titrations, the volume of $\text{NaOH}(aq)$ needed to reach the equivalence point is the same for all four titrations.	

Question 29

Skill	Learning Objective	Topic
5.F	SAP-9.A	Introduction to Acids and Bases
(A)	Incorrect. $K_w = [\text{H}^+][\text{OH}^-]$. Since one H_2O molecule ionizes to form one H^+ ion and one OH^- ion, $[\text{H}^+] = [\text{OH}^-]$, and $[\text{H}^+] = \sqrt{K_w} = \sqrt{3.0 \times 10^{-14}}$. The value $\sqrt{3.0 \times 10^{-14}}$ can be estimated to be between 1×10^{-7} and 2×10^{-7} . Since $\text{pH} = -\log[\text{H}^+]$, the pH must be slightly less than 7 (6.8).	
(B)	Correct. $K_w = [\text{H}^+][\text{OH}^-]$. Since one H_2O molecule ionizes to form one H^+ ion and one OH^- ion, $[\text{H}^+] = [\text{OH}^-]$, and $[\text{H}^+] = \sqrt{K_w} = \sqrt{3.0 \times 10^{-14}}$. The value $\sqrt{3.0 \times 10^{-14}}$ can be estimated to be between 1×10^{-7} and 2×10^{-7} and the log between -7 and -6 . Since $\text{pH} = -\log[\text{H}^+]$, the pH must be slightly less than 7 (6.8). Qualitatively, since K_w at 40°C is greater than K_w at 25°C (1×10^{-14}), this means that water ionizes more at the higher temperature, so the concentration of H^+ must be higher at the higher temperature, so the pH must be lower than 7.	
(C)	Incorrect. The pH of pure water is 7.0 at 25°C , but since the ionization constant varies with temperature, the pH of pure water also varies with temperature. $K_w = [\text{H}^+][\text{OH}^-]$. Since one H_2O molecule ionizes to form one H^+ ion and one OH^- ion, $[\text{H}^+] = [\text{OH}^-]$, and $[\text{H}^+] = \sqrt{K_w} = \sqrt{3.0 \times 10^{-14}}$. The value $\sqrt{3.0 \times 10^{-14}}$ can be estimated to be between 1×10^{-7} and 2×10^{-7} . Since $\text{pH} = -\log[\text{H}^+]$, the pH must be slightly less than 7 (6.8).	
(D)	Incorrect. $K_w = [\text{H}^+][\text{OH}^-]$. Since one H_2O molecule ionizes to form one H^+ ion and one OH^- ion, $[\text{H}^+] = [\text{OH}^-]$, and $[\text{H}^+] = \sqrt{K_w} = \sqrt{3.0 \times 10^{-14}}$. The value $\sqrt{3.0 \times 10^{-14}}$ can be estimated to be between 1×10^{-7} and 2×10^{-7} . Since $\text{pH} = -\log[\text{H}^+]$, the pH must be slightly less than 7 (6.8).	

Question 30

Skill	Learning Objective	Topic
4.A	ENE-2.C	Heat Transfer and Thermal Equilibrium
(A)	Incorrect. The average kinetic energy of a sample of gas particles is greater at a higher temperature. When the two samples are combined, the particles of the two different gases collide, and the net effect is that kinetic energy is transferred from the particles at the higher temperature to the particles at the lower temperature until the particles of the two gases have the same average kinetic energy and are at the same temperature. The Xe atoms started at the higher temperature, so the average kinetic energy of the Xe atoms decreases when the two gases are combined. Since $KE = \frac{1}{2}mv^2$ (where v is speed), if the average kinetic energy decreases then the average speed must also decrease.	
(B)	Incorrect. The average kinetic energy of a sample of gas particles is greater at a higher temperature. When the two samples are combined, the particles of the two different gases collide, and the net effect is that kinetic energy is transferred from the particles at the higher temperature to the particles at the lower temperature until the particles of the two gases have the same average kinetic energy and are at the same temperature. The Xe atoms started at the higher temperature, so the average kinetic energy of the Xe atoms decreases when the two gases are combined. Since $KE = \frac{1}{2}mv^2$ (where v is speed), if the average kinetic energy decreases then the average speed must also decrease.	
(C)	Incorrect. The average kinetic energy of a sample of gas particles is greater at a higher temperature. When the two samples are combined, the particles of the two different gases collide, and the net effect is that kinetic energy is transferred from the particles at the higher temperature to the particles at the lower temperature until the particles of the two gases have the same average kinetic energy and are at the same temperature. The Xe atoms started at the higher temperature, so the average kinetic energy of the Xe atoms decreases when the two gases are combined. Since $KE = \frac{1}{2}mv^2$ (where v is speed), if the average kinetic energy decreases then the average speed must also decrease.	
(D)	Correct. The average kinetic energy of a sample of gas particles is greater at a higher temperature. When the two samples are combined, the particles of the two different gases collide, and the net effect is that kinetic energy is transferred from the particles at the higher temperature to the particles at the lower temperature until the particles of the two gases have the same average kinetic energy and are at the same temperature. The Xe atoms started at the higher temperature, so the average kinetic energy of the Xe atoms decreases when the two gases are combined. Since $KE = \frac{1}{2}mv^2$ (where v is speed), if the average kinetic energy decreases then the average speed must also decrease.	

Question 31

Skill	Learning Objective	Topic
4.C	SAP-2.B	Valence Electrons and ionic Compounds
(A)	Incorrect. Ionization energy is the energy needed to remove an electron from an atom. When Na and Mg react with water, they lose electrons to become positive ions. The first ionization energy of Na is less than that of Mg, so Na loses an electron more easily than Mg does.	
(B)	Incorrect. Ionization energy is the energy needed to remove an electron from an atom. When Na and Mg react with water, they lose electrons to become positive ions. The first ionization energy of Na is less than that of Mg, so Na loses an electron more easily than Mg does.	
(C)	Incorrect. Ionization energy is the energy needed to remove an electron from an atom. When Na and Mg react with water, they lose electrons to become positive ions. The first ionization energy of Na is less than that of Mg, so Na loses an electron more easily than Mg does.	
(D)	Correct. Ionization energy is the energy needed to remove an electron from an atom. When Na and Mg react with water, they lose electrons to become positive ions. The first ionization energy of Na is less than that of Mg, so Na loses an electron more easily than Mg does.	

Question 32

Skill	Learning Objective	Topic
5.F	SAP-9.A	Introduction to Acids and Bases
(A)	Incorrect. The pH scale is logarithmic. For the pH to increase by 1, the concentration of the acid must decrease by a factor of 10. Therefore, the 30.0 mL sample must have been diluted to form a 300. mL sample, which is done by adding approximately 270. mL of water.	
(B)	Incorrect. The pH scale is logarithmic. For the pH to increase by 1, the concentration of the acid must decrease by a factor of 10. Therefore, the 30.0 mL sample must have been diluted to form a 300. mL sample, which is done by adding approximately 270. mL of water.	
(C)	Correct. The pH scale is logarithmic. For the pH to increase by 1, the concentration of the acid must decrease by a factor of 10. Therefore, the 30.0 mL sample must have been diluted to form a 300. mL sample, which is done by adding approximately 270. mL of water.	
(D)	Incorrect. The pH scale is logarithmic. For the pH to increase by 1, the concentration of the acid must decrease by a factor of 10. Therefore, the 30.0 mL sample must have been diluted to form a 300. mL sample, which is done by adding approximately 270. mL of water.	

Question 33

Skill	Learning Objective	Topic
5.B	ENE-4.C	Gibbs Free-Energy and Thermodynamic Favorability
(A)	Incorrect. When the equilibrium constant is greater than 1.0, ΔG° is less than 0 (the reaction is thermodynamically favorable), and when the equilibrium constant is less than 1.0, ΔG° is greater than 0 (the reaction is not thermodynamically favorable). Since $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$, $\Delta H^\circ < 0$ and $\Delta S^\circ < 0$ results in a negative ΔG° at low temperatures and a positive ΔG° at high temperatures.	
(B)	Incorrect. When the equilibrium constant is greater than 1.0, ΔG° is less than 0 (the reaction is thermodynamically favorable), and when the equilibrium constant is less than 1.0, ΔG° is greater than 0 (the reaction is not thermodynamically favorable). Since $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$, $\Delta H^\circ < 0$ and $\Delta S^\circ < 0$ results in a negative ΔG° at low temperatures and a positive ΔG° at high temperatures.	
(C)	Incorrect. When the equilibrium constant is greater than 1.0, ΔG° is less than 0 (the reaction is thermodynamically favorable), and when the equilibrium constant is less than 1.0, ΔG° is greater than 0 (the reaction is not thermodynamically favorable). Since $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$, $\Delta H^\circ < 0$ and $\Delta S^\circ < 0$ results in a negative ΔG° at low temperatures and a positive ΔG° at high temperatures.	
(D)	Correct. When the equilibrium constant is greater than 1.0, ΔG° is less than 0 (the reaction is thermodynamically favorable), and when the equilibrium constant is less than 1.0, ΔG° is greater than 0 (the reaction is not thermodynamically favorable). Since $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$, $\Delta H^\circ < 0$, and $\Delta S^\circ < 0$ results in a negative ΔG° at low temperatures and a positive ΔG° at high temperatures.	

Question 34

Skill	Learning Objective	Topic
6.F	SPQ-5.A	Introduction to Solubility Equilibria
(A)	Incorrect. The concentration of ions in a saturated solution of an ionic solid at a given temperature is constant. As water evaporates from the solution, enough of the solute precipitates to keep the concentration of the ions in solution constant.	
(B)	Incorrect. The concentration of ions in a saturated solution of an ionic solid at a given temperature is constant. As water evaporates from the solution, enough of the solute precipitates to keep the concentration of the ions in solution constant.	
(C)	Incorrect. The concentration of ions in a saturated solution of an ionic solid at a given temperature is constant. As water evaporates from the solution, enough of the solute precipitates to keep the concentration of the ions in solution constant.	
(D)	Correct. The concentration of ions in a saturated solution of an ionic solid at a given temperature is constant. As water evaporates from the solution, enough of the solute precipitates to keep the concentration of the ions in solution constant.	

Question 35

Skill	Learning Objective	Topic
5.E	TRA-1.B	Net Ionic Equations
(A)	Incorrect. Since the question asks for the species that react and the species that are produced, the equation should be written in net ionic form. The products of the reaction are $\text{Ca}^{2+}(aq)$, $\text{CO}_2(g)$, and $\text{H}_2\text{O}(l)$.	
(B)	Correct. Since the question asks for the species that react and the species that are produced, the equation should be written in net ionic form. The products of the reaction are $\text{Ca}^{2+}(aq)$, $\text{CO}_2(g)$, and $\text{H}_2\text{O}(l)$.	
(C)	Incorrect. Since the question asks for the species that react and the species that are produced, the equation should be written in net ionic form. The products of the reaction are $\text{Ca}^{2+}(aq)$, $\text{CO}_2(g)$, and $\text{H}_2\text{O}(l)$.	
(D)	Incorrect. Since the question asks for the species that react and the species that are produced, the equation should be written in net ionic form. The products of the reaction are $\text{Ca}^{2+}(aq)$, $\text{CO}_2(g)$, and $\text{H}_2\text{O}(l)$. $\text{Cl}^-(aq)$ is not included because it is a spectator ion.	

Question 36

Skill	Learning Objective	Topic
5.F	SPQ-2.B	Composition of Mixtures
(A)	Incorrect. The difference in pressure is $0.870 \text{ atm} - 0.800 \text{ atm} = 0.070 \text{ atm}$. According to the graph, this pressure corresponds to a mass of $\text{CaCO}_3(s)$ of 0.145 g. The mass of eggshell used was 0.200 g. The mass percent of $\text{CaCO}_3(s)$ in the eggshell was thus approximately $\frac{0.145 \text{ g}}{0.200 \text{ g}} \times 100 = 73\%$.	
(B)	Incorrect. The difference in pressure is $0.870 \text{ atm} - 0.800 \text{ atm} = 0.070 \text{ atm}$. According to the graph, this pressure corresponds to a mass of $\text{CaCO}_3(s)$ of 0.145 g. The mass of eggshell used was 0.200 g. The mass percent of $\text{CaCO}_3(s)$ in the eggshell was thus approximately $\frac{0.145 \text{ g}}{0.200 \text{ g}} \times 100 = 73\%$.	
(C)	Incorrect. The difference in pressure is $0.870 \text{ atm} - 0.800 \text{ atm} = 0.070 \text{ atm}$. According to the graph, this pressure corresponds to a mass of $\text{CaCO}_3(s)$ of 0.145 g. The mass of eggshell used was 0.200 g. The mass percent of $\text{CaCO}_3(s)$ in the eggshell was thus approximately $\frac{0.145 \text{ g}}{0.200 \text{ g}} \times 100 = 73\%$.	
(D)	Correct. The difference in pressure is $0.870 \text{ atm} - 0.800 \text{ atm} = 0.070 \text{ atm}$. According to the graph, this pressure corresponds to a mass of $\text{CaCO}_3(s)$ of 0.145 g. The mass of eggshell used was 0.200 g. The mass percent of $\text{CaCO}_3(s)$ in the eggshell was thus approximately $\frac{0.145 \text{ g}}{0.200 \text{ g}} \times 100 = 73\%$.	

Question 37

Skill	Learning Objective	Topic
5.F	SPQ-3.A	Solutions and Mixtures
(A)	Incorrect. According to the graph, 0.095 atm of gas corresponds to approximately 0.20 g of CaCO_3. Since the molar mass of CaCO_3 is 100. g/mol, this corresponds to 0.0020 mol of CaCO_3. Each mole of CaCO_3 requires two moles of H^+ ions in the reaction, so 0.0040 mol of acid is needed. $\frac{0.0040 \text{ mol}}{4.0 \text{ mL}} = \frac{x \text{ mol}}{1000 \text{ mL}}$, so $x = 1.0$, and the concentration of $\text{HCl}(aq)$ was 1.0 M.	
(B)	Incorrect. According to the graph, 0.095 atm of gas corresponds to approximately 0.20 g of CaCO_3. Since the molar mass of CaCO_3 is 100. g/mol, this corresponds to 0.0020 mol of CaCO_3. Each mole of CaCO_3 requires two moles of H^+ ions in the reaction, so 0.0040 mol of acid is needed. $\frac{0.0040 \text{ mol}}{4.0 \text{ mL}} = \frac{x \text{ mol}}{1000 \text{ mL}}$, so $x = 1.0$, and the concentration of $\text{HCl}(aq)$ was 1.0 M.	
(C)	Incorrect. According to the graph, 0.095 atm of gas corresponds to approximately 0.20 g of CaCO_3. Since the molar mass of CaCO_3 is 100. g/mol, this corresponds to 0.0020 mol of CaCO_3. Each mole of CaCO_3 requires two moles of H^+ ions in the reaction, so 0.0040 mol of acid is needed. $\frac{0.0040 \text{ mol}}{4.0 \text{ mL}} = \frac{x \text{ mol}}{1000 \text{ mL}}$, so $x = 1.0$, and the concentration of $\text{HCl}(aq)$ was 1.0 M.	
(D)	Correct. According to the graph, 0.095 atm of gas corresponds to approximately 0.20 g of CaCO_3. Since the molar mass of CaCO_3 is 100. g/mol, this corresponds to 0.0020 mol of CaCO_3. Each mole of CaCO_3 requires two moles of H^+ ions in the reaction, so 0.0040 mol of acid is needed. $\frac{0.0040 \text{ mol}}{4.0 \text{ mL}} = \frac{x \text{ mol}}{1000 \text{ mL}}$, so $x = 1.0$, and the concentration of $\text{HCl}(aq)$ was 1.0 M.	

Question 38

Skill	Learning Objective	Topic
1.B	TRA-3.A	Reaction Rates
(A)	Incorrect. Using a weak acid instead of a strong acid would not increase the rate of the reaction. Increasing the surface area of the reactant increases the rate of the reaction, because more of the solid CaCO_3 is in contact with the H^+ ions in solution.	
(B)	Incorrect. Reaction rates typically increase when the temperature is increased, not when it is decreased. Increasing the surface area of the reactant increases the rate of the reaction, because more of the solid CaCO_3 is in contact with the H^+ ions in solution.	
(C)	Incorrect. Reducing the volume of the reaction vessel would not increase the rate of the reaction—it would not increase concentrations in an aqueous solution. Increasing the surface area of the reactant increases the rate of the reaction, because more of the solid CaCO_3 is in contact with the H^+ ions in solution.	
(D)	Correct. Increasing the surface area of the reactant increases the rate of the reaction, because more of the solid CaCO_3 is in contact with the H^+ ions in solution.	

Question 39

Skill	Learning Objective	Topic
5.E	SAP-9.C	Weak Acid and Base Equilibria
(A)	Incorrect. The Na^+ ion does not react with water as shown in the equation. $\text{NH}_4^+(aq)$ partially hydrolyzes (reacts with water) to form $\text{NH}_3(aq)$ and $\text{H}_3\text{O}^+(aq)$. Since $\text{H}_3\text{O}^+(aq)$ is a product, the resulting solution is slightly acidic.	
(B)	Incorrect. The solution formed in this reaction is slightly basic (OH^- ions are formed). $\text{NH}_4^+(aq)$ partially hydrolyzes (reacts with water) to form $\text{NH}_3(aq)$ and $\text{H}_3\text{O}^+(aq)$. Since $\text{H}_3\text{O}^+(aq)$ is a product, the resulting solution is slightly acidic.	
(C)	Correct. $\text{NH}_4^+(aq)$ partially hydrolyzes (reacts with water) to form $\text{NH}_3(aq)$ and $\text{H}_3\text{O}^+(aq)$. Since $\text{H}_3\text{O}^+(aq)$ is a product, the resulting solution is slightly acidic.	
(D)	Incorrect. The Cl^- ion does not react with water as shown in the equation. $\text{NH}_4^+(aq)$ partially hydrolyzes (reacts with water) to form $\text{NH}_3(aq)$ and $\text{H}_3\text{O}^+(aq)$. Since $\text{H}_3\text{O}^+(aq)$ is a product, the resulting solution is slightly acidic.	

Question 40

Skill	Learning Objective	Topic
6.E	ENE-4.D	Thermodynamic and Kinetic Control
(A)	Incorrect. The value of ΔG° does not indicate anything about the rate of a reaction, just its thermodynamic favorability. Reactions with high activation energies are very slow. Energy is needed to break bonds in reactant molecules, and diamond is a covalent network solid with strong bonds in four different directions.	
(B)	Incorrect. The change in entropy does not affect the rate of the reaction. Reactions with high activation energies are very slow. Energy is needed to break bonds in reactant molecules, and diamond is a covalent network solid with strong bonds in four different directions.	
(C)	Correct. Reactions with high activation energies are very slow. Energy is needed to break bonds in reactant molecules, and diamond is a covalent network solid with strong bonds in four different directions.	
(D)	Incorrect. The value of ΔG° is negative, so the reaction is thermodynamically favorable. Reactions with high activation energies are very slow. Energy is needed to break bonds in reactant molecules, and diamond is a covalent network solid with strong bonds in four different directions.	

Question 41

Skill	Learning Objective	Topic
4.A	SAP-9.E	Acid-Base Titrations
(A)	Incorrect. At the equivalence point, the number of moles of $\text{OH}^- (aq)$ added is equal to the initial number of moles of $\text{C}_9\text{H}_8\text{O}_4(aq)$, and the reaction shown has gone essentially to completion. Therefore, the concentration of $\text{C}_9\text{H}_7\text{O}_4^- (aq)$ is much greater than that of $\text{C}_9\text{H}_8\text{O}_4(aq)$.	
(B)	Correct. At the equivalence point, the number of moles of $\text{OH}^- (aq)$ added is equal to the initial number of moles of $\text{C}_9\text{H}_8\text{O}_4(aq)$, and the reaction shown has gone essentially to completion. Therefore, the concentration of $\text{C}_9\text{H}_7\text{O}_4^- (aq)$ is much greater than that of $\text{C}_9\text{H}_8\text{O}_4(aq)$.	
(C)	Incorrect. At the equivalence point, the number of moles of $\text{OH}^- (aq)$ added is equal to the initial number of moles of $\text{C}_9\text{H}_8\text{O}_4(aq)$, and the reaction shown has gone essentially to completion. Therefore, the concentration of $\text{C}_9\text{H}_7\text{O}_4^- (aq)$ is much greater than that of $\text{C}_9\text{H}_8\text{O}_4(aq)$.	
(D)	Incorrect. At the equivalence point, the number of moles of $\text{OH}^- (aq)$ added is equal to the initial number of moles of $\text{C}_9\text{H}_8\text{O}_4(aq)$, and the reaction shown has gone essentially to completion. Therefore, the concentration of $\text{C}_9\text{H}_7\text{O}_4^- (aq)$ is much greater than that of $\text{C}_9\text{H}_8\text{O}_4(aq)$.	

Question 42

Skill	Learning Objective	Topic
1.A	SAP-4.C	VSEPR and Bond Hybridization
(A)	Incorrect. The oxygen atom in the $\text{H} - \text{O} - \text{C}$ bond has two lone pairs of electrons, so the geometry of the electron domains around the oxygen atom is roughly tetrahedral. The bond angle is thus close to 109° .	
(B)	Incorrect. The oxygen atom in the $\text{H} - \text{O} - \text{C}$ bond has two lone pairs of electrons, so the geometry of the electron domains around the oxygen atom is roughly tetrahedral. The bond angle is thus close to 109° .	
(C)	Correct. The oxygen atom in the $\text{H} - \text{O} - \text{C}$ bond has two lone pairs of electrons, so the geometry of the electron domains around the oxygen atom is roughly tetrahedral. The bond angle is thus close to 109° .	
(D)	Incorrect. The oxygen atom in the $\text{H} - \text{O} - \text{C}$ bond has two lone pairs of electrons, so the geometry of the electron domains around the oxygen atom is roughly tetrahedral. The bond angle is thus close to 109° .	

Question 43

Skill	Learning Objective	Topic
1.A	SAP-4.C	VSEPR and Bond Hybridization
(A)	Incorrect. The leftmost carbon atom is bonded to four atoms and it does not have any nonbonding pairs of electrons, so it has four electron domains. The geometry of the electron domains around the leftmost carbon atom is thus roughly tetrahedral.	
(B)	Correct. The carbon atom bonded to two oxygen atoms and another carbon atom does not have any nonbonding pairs of electrons, thus the central carbon atom has three electron domains. According to VSEPR theory, all four atoms are in the same plane (sp^2 hybridization).	
(C)	Incorrect. The leftmost carbon atom is bonded to four atoms and it does not have any nonbonding pairs of electrons on it, so it has four electron domains. The geometry of the electron domains around the leftmost carbon atom is thus roughly tetrahedral and the N – C – C bond angle is roughly 109°.	
(D)	Incorrect. The nitrogen atom is bonded to four atoms and it does not have any nonbonding pairs of electrons on it, so it has four electron domains. The geometry of the electron domains around the nitrogen atom is thus roughly tetrahedral.	

Question 44

Skill	Learning Objective	Topic
1.A	SAP-5.B	Properties of Solids
(A)	Correct. The sodium ion and the glycinate ion form the ionic compound sodium glycinate.	
(B)	Incorrect. The sodium ion and the glycinate ion form the ionic compound sodium glycinate, which, like all sodium salts, is soluble in water.	
(C)	Incorrect. The sodium ion and the glycinate ion form the ionic compound sodium glycinate, which does not evaporate easily.	
(D)	Incorrect. The sodium ion and the glycinate ion form the ionic compound sodium glycinate. The glycinate ion is formed through the loss of a proton (H^+) and would not lose another proton in aqueous solution to form an acidic solution.	

Question 45

Skill	Learning Objective	Topic
1.A	SAP-3.A	Types of Chemical Bonds
(A)	Incorrect. Polar covalent bonds are the <u>intramolecular</u> forces within a carbon tetrachloride molecule. Carbon tetrachloride molecules have equivalent bonds that are symmetrically arranged around the carbon atom; thus, the bond dipoles cancel and the molecules are nonpolar. The attractions between molecules are London dispersion forces, which result from temporary dipoles.	
(B)	Incorrect. Polar (not nonpolar) covalent bonds are the <u>intramolecular</u> forces within a carbon tetrachloride molecule. Carbon tetrachloride molecules have equivalent bonds that are symmetrically arranged around the carbon atom; thus, the bond dipoles cancel and the molecules are nonpolar. The attractions between molecules are London dispersion forces, which result from temporary dipoles.	
(C)	Correct. Carbon tetrachloride molecules have equivalent bonds that are symmetrically arranged around the carbon atom; thus, the bond dipoles cancel and the molecules are nonpolar. The attractions between molecules are London dispersion forces, which result from temporary dipoles.	
(D)	Incorrect. Carbon tetrachloride molecules have equivalent bonds that are symmetrically arranged around the carbon atom; thus, the bond dipoles cancel and the molecules are nonpolar. The attractions between molecules are London dispersion forces, which result from temporary dipoles.	

Question 46

Skill	Learning Objective	Topic
2.C	ENE-2.D	Heat Capacity and Calorimetry
(A)	Incorrect. The mass of the water is also needed.	
(B)	Incorrect. The mass of the $\text{KCl}(s)$ is also needed.	
(C)	Correct. The formula $q = mc\Delta T$ is the formula used to determine the amount of heat (q) released or absorbed when $\text{KCl}(s)$ is dissolved in water. The experimental procedure is to dissolve a known amount of $\text{KCl}(s)$ in a known mass of water and to measure the temperature before and after the dissolution. The enthalpy change for the dissolution of one mole of $\text{KCl}(s)$ can then be calculated.	
(D)	Incorrect. The atmospheric pressure is not needed, as would be the case if the vapor pressure of the water was needed.	

Question 47

Skill	Learning Objective	Topic
5.F	TRA-8.B	Reaction Quotient and Le Châtelier's Principle
(A)	Correct. At these initial concentrations, $Q = \frac{[XY]^2}{[X_2][Y_2]} = \frac{(0.20)^2}{(0.40)(0.40)} = \frac{0.040}{0.16} = 0.25.$ Since $Q < K_c$, the system will shift to the right and more product will form.	
(B)	Incorrect. At these initial concentrations, $Q = \frac{[XY]^2}{[X_2][Y_2]} = \frac{(0.90)(0.90)}{(0.30)(0.30)} = \frac{(3)(3)}{(1)(1)} = 9.0.$ Since $Q > K_c$, the system will shift to the left and more reactants will form.	
(C)	Incorrect. At these initial concentrations, $Q = \frac{[XY]^2}{[X_2][Y_2]} = \frac{(0.30)(0.30)}{(0.15)(0.15)} = \frac{(2)(2)}{(1)(1)} = 4.0.$ Since $Q > K_c$, the system will shift to the left and more reactants will form.	
(D)	Incorrect. At these initial concentrations, $Q = \frac{[XY]^2}{[X_2][Y_2]} = \frac{(0.20)(0.20)}{(0.10)(0.10)} = \frac{(2)(2)}{(1)(1)} = 4.0.$ Since $Q > K_c$, the system will shift to the left and more reactants will form.	

Question 48

Skill	Learning Objective	Topic
6.E	SAP-5.A	Intermolecular Forces
(A)	Incorrect. The bond dipoles cancel in the <i>trans</i> -isomer because they are equal and pointed in opposite directions. The bond dipoles in the <i>cis</i> -isomer do not cancel, so the molecules have a net dipole moment, which means that the <i>cis</i> -isomer has dipole-dipole forces in addition to London dispersion forces. Since the <i>trans</i> -isomer has only London dispersion forces (which are equal in strength to the London dispersion forces in the <i>cis</i> -isomer), it is easier for the molecules of the <i>trans</i> -isomer to escape the surface of the liquid; thus, it has a higher equilibrium vapor pressure.	
(B)	Incorrect. The bond dipoles cancel in the <i>trans</i> -isomer because they are equal and pointed in opposite directions. The bond dipoles in the <i>cis</i> -isomer do not cancel, so the molecules have a net dipole moment, which means that the <i>cis</i> -isomer has dipole-dipole forces in addition to London dispersion forces. Since the <i>trans</i> -isomer has only London dispersion forces (which are equal in strength to the London dispersion forces in the <i>cis</i> -isomer), it is easier for the molecules of the <i>trans</i> -isomer to escape the surface of the liquid; thus, it has a higher equilibrium vapor pressure.	
(C)	Incorrect. The bond dipoles cancel in the <i>trans</i> -isomer because they are equal and pointed in opposite directions. The bond dipoles in the <i>cis</i> -isomer do not cancel, so the molecules have a net dipole moment, which means that the <i>cis</i> -isomer has dipole-dipole forces in addition to London dispersion forces. Since the <i>trans</i> -isomer has only London dispersion forces (which are equal in strength to the London dispersion forces in the <i>cis</i> -isomer), it is easier for the molecules of the <i>trans</i> -isomer to escape the surface of the liquid; thus, it has a higher equilibrium vapor pressure.	
(D)	Correct. The bond dipoles cancel in the <i>trans</i> -isomer because they are equal and pointed in opposite directions. The bond dipoles in the <i>cis</i> -isomer do not cancel, so the molecules have a net dipole moment, which means that the <i>cis</i> -isomer has dipole-dipole forces in addition to London dispersion forces. Since the <i>trans</i> -isomer has only London dispersion forces (which are equal in strength to the London dispersion forces in the <i>cis</i> -isomer), it is easier for the molecules of the <i>trans</i> -isomer to escape the surface of the liquid; thus, it has a higher equilibrium vapor pressure.	

Question 49

Skill	Learning Objective	Topic
5.F	SPQ-5.B	Common-Ion Effect
(A)	Incorrect. $K_{sp} = [\text{Ca}^{2+}][\text{F}^-]^2$, so $4.0 \times 10^{-11} = [\text{Ca}^{2+}][4.0 \times 10^{-5}]^2$, and $[\text{Ca}^{2+}] = \frac{4.0 \times 10^{-11}}{16 \times 10^{-10}} = \frac{4.0 \times 10^{-11}}{1.6 \times 10^{-9}} = 2.5 \times 10^{-11-(-9)} = 0.025 \text{ M}.$	
(B)	Correct. $K_{sp} = [\text{Ca}^{2+}][\text{F}^-]^2$, so $4.0 \times 10^{-11} = [\text{Ca}^{2+}][4.0 \times 10^{-5}]^2$, and $[\text{Ca}^{2+}] = \frac{4.0 \times 10^{-11}}{16 \times 10^{-10}} = \frac{4.0 \times 10^{-11}}{1.6 \times 10^{-9}} = 2.5 \times 10^{-11-(-9)} = 0.025 \text{ M}.$	
(C)	Incorrect. $K_{sp} = [\text{Ca}^{2+}][\text{F}^-]^2$, so $4.0 \times 10^{-11} = [\text{Ca}^{2+}][4.0 \times 10^{-5}]^2$, and $[\text{Ca}^{2+}] = \frac{4.0 \times 10^{-11}}{16 \times 10^{-10}} = \frac{4.0 \times 10^{-11}}{1.6 \times 10^{-9}} = 2.5 \times 10^{-11-(-9)} = 0.025 \text{ M}.$	
(D)	Incorrect. $K_{sp} = [\text{Ca}^{2+}][\text{F}^-]^2$, so $4.0 \times 10^{-11} = [\text{Ca}^{2+}][4.0 \times 10^{-5}]^2$, and $[\text{Ca}^{2+}] = \frac{4.0 \times 10^{-11}}{16 \times 10^{-10}} = \frac{4.0 \times 10^{-11}}{1.6 \times 10^{-9}} = 2.5 \times 10^{-11-(-9)} = 0.025 \text{ M}.$	


Question 50

Skill	Learning Objective	Topic
5.F	TRA-7.D	Properties of the Equilibrium Constant
(A)	Incorrect. The third equation is the sum of the first equation and the reverse of the second equation. This means that K_{eq} of the third equation is equal to K_{eq} of the first equation multiplied by $\frac{1}{K_{eq}}$ of the second equation, or $(1 \times 10^{-6}) \times \frac{1}{(1 \times 10^{-32})} = (1 \times 10^{-6})(1 \times 10^{32}) = 1 \times 10^{26}.$ Since K_{eq} is greater than 1, ΔG° must be negative.	
(B)	Incorrect. The third equation is the sum of the first equation and the reverse of the second equation. This means that K_{eq} of the third equation is equal to K_{eq} of the first equation multiplied by $\frac{1}{K_{eq}}$ of the second equation, or $(1 \times 10^{-6}) \times \frac{1}{(1 \times 10^{-32})} = (1 \times 10^{-6})(1 \times 10^{32}) = 1 \times 10^{26}.$ Since K_{eq} is greater than 1, ΔG° must be negative.	
(C)	Incorrect. The third equation is the sum of the first equation and the reverse of the second equation. This means that K_{eq} of the third equation is equal to K_{eq} of the first equation multiplied by $\frac{1}{K_{eq}}$ of the second equation, or $(1 \times 10^{-6}) \times \frac{1}{(1 \times 10^{-32})} = (1 \times 10^{-6})(1 \times 10^{32}) = 1 \times 10^{26}.$ Since K_{eq} is greater than 1, ΔG° must be negative.	
(D)	Correct. The third equation is the sum of the first equation and the reverse of the second equation. This means that K_{eq} of the third equation is equal to K_{eq} of the first equation multiplied by $\frac{1}{K_{eq}}$ of the second equation, or $(1 \times 10^{-6}) \times \frac{1}{(1 \times 10^{-32})} = (1 \times 10^{-6})(1 \times 10^{32}) = 1 \times 10^{26}.$ Since K_{eq} is greater than 1, ΔG° must be negative.	

AP[®] CHEMISTRY

2019 SCORING GUIDELINES

Question 1



Ethanol, $\text{C}_2\text{H}_5\text{OH}$, will combust in air according to the equation above.

- (a) Is $\text{O}_2(g)$ oxidized in the reaction, or is it reduced? Justify your answer in terms of oxidation numbers.

O_2 is reduced; its oxidation number changes from 0 in O_2 to -2 in CO_2 and H_2O .	1 point is earned for the correct answer with an appropriate justification.
--	---

- (b) When a sample of $\text{C}_2\text{H}_5\text{OH}$ was combusted, the volume of $\text{CO}_2(g)$ produced was 18.0 L when measured at 21.7°C and 1.03 atm. Determine the number of moles of $\text{CO}_2(g)$ that was produced.

$n = \frac{PV}{RT} = \frac{(1.03 \text{ atm})(18.0 \text{ L})}{(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(273.15 + 21.7) \text{ K}} = 0.766 \text{ mol CO}_2$	1 point is earned for the correct calculation.
--	--

- (c) Determine the volume of $\text{C}_2\text{H}_5\text{OH}(l)$, in mL, that was combusted to produce the volume of $\text{CO}_2(g)$ collected in part (b). (The density of $\text{C}_2\text{H}_5\text{OH}(l)$ is 0.79 g/mL .)

$0.766 \text{ mol CO}_2 \times \frac{1 \text{ mol C}_2\text{H}_5\text{OH}}{2 \text{ mol CO}_2} \times \frac{46.07 \text{ g C}_2\text{H}_5\text{OH}}{1 \text{ mol C}_2\text{H}_5\text{OH}} \times \frac{1 \text{ mL}}{0.79 \text{ g}} = 22 \text{ mL}$	1 point is earned for the correct stoichiometry for moles of $\text{C}_2\text{H}_5\text{OH}$ and CO_2 . 1 point is earned for the correct volume of $\text{C}_2\text{H}_5\text{OH}$.
---	---

- (d) Determine the amount of heat, in kJ, that is released by the combustion reaction.

$q = n \Delta H_{\text{comb}}^\circ$ $q = 0.766 \text{ mol CO}_2 \times \frac{1 \text{ mol}_{\text{rxn}}}{2 \text{ mol CO}_2} \times \frac{-1270 \text{ kJ}}{1 \text{ mol}_{\text{rxn}}} = -486 \text{ kJ}$ The amount of heat released is 486 kJ.	1 point is earned for a correct calculation (negative sign is not required).
---	--

The combustion reaction occurred in a closed room containing $5.56 \times 10^4 \text{ g}$ of air originally at 21.7°C . Assume that all of the heat produced by the reaction was absorbed by the air (specific heat = $1.005 \text{ J/(g } ^\circ\text{C)}$) in the room.

AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 1 (continued)

(e) Determine the final temperature of the air in the room after the combustion.

$q_{air} = -q_{rxn} = +486 \text{ kJ} = +486,000 \text{ J}$ $q = mc\Delta T \Rightarrow \Delta T = \frac{q}{mc} = \frac{486,000 \text{ J}}{(5.56 \times 10^4 \text{ g})(1.005 \text{ J/(g } ^\circ\text{C)})} = 8.70^\circ\text{C}$ $T_{final} = 21.7^\circ\text{C} + 8.70^\circ\text{C} = 30.4^\circ\text{C}$	1 point is earned for the correct final temperature.
--	--

(f) The boiling points of ethanol and dimethyl ether are provided in the following table. Identify the intermolecular force that is most responsible for the difference in boiling point. In terms of this force, explain why the boiling point of ethanol is higher than that of dimethyl ether.

Compound	Structural Formula	Boiling Point ($^\circ\text{C}$)
Ethanol	<pre> H H H — C — C — O — H H H </pre>	78
Dimethyl ether	<pre> H H H — C — O — C — H H H </pre>	–24

Hydrogen bonding. Ethanol can form hydrogen bonds between its molecules, whereas dimethyl ether cannot. The attractions between molecules are stronger in ethanol than in dimethyl ether; therefore, the boiling point of ethanol is higher.	1 point is earned for identifying hydrogen bonding as the relevant force. 1 point is earned for linking the greater strength of the intermolecular forces to a higher boiling point.
--	--

One method of producing ethanol is by the reaction of ethene and water, as represented below.



AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 1 (continued)

- (g) A chemist wants to run the reaction and maximize the amount of $\text{C}_2\text{H}_5\text{OH(g)}$ produced. Identify two ways the chemist could change the reaction conditions (other than adding or removing any chemical species) to favor the formation of more product. Justify your answer.


Decrease the volume of the reaction container. There are fewer moles of gaseous product than moles of gaseous reactants; therefore, an increase in pressure due to a decrease in volume would favor the formation of product.

Lower the temperature. A lower temperature favors an exothermic reaction, leading to the conversion of reactants into product.

1 point is earned for EACH valid answer with justification (2 points total).


AP[®] CHEMISTRY
2019 SCORING GUIDELINES


Question 2




Hydrogen peroxide, H_2O_2 , decomposes according to the equation above. This reaction is thermodynamically favorable at room temperature.

- (a) A particulate representation of the reactants is shown below in the box on the left. In the box below on the right, draw the particulate representation of all the molecules that would be produced from these four reactant molecules.



	1 point is earned for correctly representing products as H_2O and O_2 molecules. 1 point is earned for atom balance.
--	---

- (b) Shown below is a potential energy diagram for the uncatalyzed decomposition of $\text{H}_2\text{O}_2(aq)$.



- (i) According to the diagram, is the decomposition reaction exothermic or endothermic? Justify your answer.


This reaction is exothermic because the products have lower potential energy than the reactants. Thus, the system has lost potential energy, releasing heat.

1 point is earned for indication that the reaction is exothermic, with a valid justification.


AP[®] CHEMISTRY 2019 SCORING GUIDELINES

Question 2 (continued)

- (ii) Manganese dioxide, $\text{MnO}_2(s)$, is an insoluble substance that acts as a catalyst for the decomposition reaction. On the diagram above, draw a curve to represent the reaction as it occurs in the presence of $\text{MnO}_2(s)$.



A student investigates the decomposition reaction in the laboratory. The student prepares two small beakers, adding 20.0 mL of 9.77 M $\text{H}_2\text{O}_2(aq)$ to each one. Each beaker is placed on an electronic balance. The student adds 0.10 g of $\text{MnO}_2(s)$ to the second beaker and records the mass of each beaker and its contents at 10-second intervals for one minute. The beakers and the data are shown below.



Time (seconds)	Mass of Beaker 1 + $\text{H}_2\text{O}_2(aq)$ (grams)	Mass of Beaker 2 + $\text{H}_2\text{O}_2(aq)$ + $\text{MnO}_2(s)$ (grams)
0	43.09	43.19
10	43.09	43.06
20	43.09	42.94
30	43.09	42.83
40	43.09	42.73
50	43.09	42.65
60	43.09	42.58

AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 2 (continued)

(c) For beaker 2 during the 60-second period, calculate the following.

(i) The number of moles of $\text{O}_2(\text{g})$ that was produced

$\text{mass O}_2 = 43.19 \text{ g} - 42.58 \text{ g} = 0.61 \text{ g}$ $\text{moles O}_2 = 0.61 \text{ g O}_2 \times \frac{1 \text{ mol O}_2}{32.00 \text{ g}} = 0.019 \text{ mol O}_2$	1 point is earned for the correct answer.
---	---

(ii) The mass of $\text{H}_2\text{O}_2(\text{aq})$ that decomposed

$\text{mass H}_2\text{O}_2 = 0.019 \text{ mol O}_2 \times \frac{2 \text{ mol H}_2\text{O}_2}{1 \text{ mol O}_2} \times \frac{34.02 \text{ g H}_2\text{O}_2}{1 \text{ mol H}_2\text{O}_2} = 1.3 \text{ g H}_2\text{O}_2$	1 point is earned for the correct answer.
---	---

(d) The student continues the experiment for an additional minute. For beaker 2, will the mass of $\text{H}_2\text{O}_2(\text{aq})$ consumed during the second minute be greater than, less than, or equal to the mass of $\text{H}_2\text{O}_2(\text{aq})$ consumed during the first minute? Explain your answer referring to the data in the table.

Less than. The decrease in mass becomes smaller over every subsequent 10-second interval. This decreasing trend would continue in the second minute.	1 point is earned for the correct choice with a valid explanation.
---	--

(e) Based on the data, the student claims that the catalyzed reaction has zeroth-order kinetics. Do you agree with the student's claim? Justify your answer.

Disagree. If the reaction were zeroth-order, the rate of mass loss (which corresponds to the rate of reaction) would be constant until the reactant was consumed. However, the data show that the rate of mass loss from beaker 2 is not constant.	1 point is earned for disagreement and a valid justification.
--	---

(f) A second student did the experiment using larger volumes of $9.77 \text{ M H}_2\text{O}_2(\text{aq})$. The student noticed that the reaction in beaker 2 proceeded extremely rapidly, causing some of the liquid to splash out of the beaker onto the lab table. The student claims that as a result of the loss of the liquid from the beaker, the calculated number of moles of $\text{O}_2(\text{g})$ produced is greater than the actual number of moles of $\text{O}_2(\text{g})$ produced during the first 60 seconds. Do you agree with the student? Justify your answer.

Agree. The calculated amount of O_2 produced is based on the change in mass, and in this case some of the change in mass is due to loss of solution rather than production of O_2 .	1 point is earned for agreement and a valid justification.
---	--

AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 2 (continued)

- (g) The hydrogen peroxide used in this experiment can be prepared by the reaction of solid ammonium peroxydisulfate, $(\text{NH}_4)_2\text{S}_2\text{O}_8$, with water. The products are hydrogen peroxide (H_2O_2) and ammonium bisulfate (NH_4HSO_4). Write the balanced equation for the reaction.


$(\text{NH}_4)_2\text{S}_2\text{O}_8 + 2 \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}_2 + 2 \text{NH}_4\text{HSO}_4$ OR $(\text{NH}_4)_2\text{S}_2\text{O}_8 + 2 \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}_2 + 2 \text{NH}_4^+ + 2 \text{HSO}_4^-$ OR $\text{S}_2\text{O}_8^{2-} + 2 \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}_2 + 2 \text{HSO}_4^-$	1 point is earned for a correct balanced equation.
---	--


AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 3

Sulfur trioxide, SO_3 , is commercially produced on a massive scale for use in manufacturing. It is also produced as a by-product of burning fossil fuels and is an air pollutant involved in the formation of acid rain.

- (a) In the box below, complete one of the valid Lewis electron-dot diagrams for the SO_3 molecule. Be sure to include all valence electrons, including lone pairs.




 (OR equivalent resonance form)	1 point is earned for any valid Lewis electron-dot diagram.
--	--

- (b) Predict the shape of the SO_3 molecule based on the diagram you completed above.

Trigonal planar	1 point is earned for the shape that is consistent with the answer to part (a).
------------------------	--

Under some conditions, $\text{SO}_3(g)$ decomposes to produce $\text{SO}_2(g)$ and $\text{O}_2(g)$, as represented by the equation below.




AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 3 (continued)

- (c) Under what temperature conditions will the reaction be thermodynamically favored: high temperatures, low temperatures, all temperatures, or no temperatures? Justify your answer in terms of the change in enthalpy and the change in entropy during the reaction.

The reaction will be thermodynamically favored only at high temperatures. The reaction will be thermodynamically favored when ΔG° is negative ($\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$). ΔH° for this reaction is positive, so at low temperatures (when the value of $T\Delta S^\circ$ is small), ΔG° is positive and the reaction is not thermodynamically favored. Because ΔS° for this reaction is positive, high-temperature conditions give a large positive value for $T\Delta S^\circ$. This will result in a negative ΔG°, and the reaction will be thermodynamically favored.	1 point is earned for correctly linking ΔH° to unfavorability at low temperatures. 1 point is earned for assigning the correct sign to ΔS° and linking it to favorability at high temperatures.
---	---

A chemist fills a rigid vessel with $\text{SO}_3(\text{g})$ at a certain temperature until the pressure of $\text{SO}_3(\text{g})$ in the container is 0.83 atm. The $\text{SO}_3(\text{g})$ decomposes as the partial pressures of $\text{SO}_3(\text{g})$, $\text{SO}_2(\text{g})$, and $\text{O}_2(\text{g})$ in the container are monitored over time, as shown in the following graph. A change is made to the system at time t_1 .



- (d) Answer the following based on the graph.

(i) Draw an X on the time axis to show when the system first reached equilibrium.

See graph above. The drawing should show an X on the time axis approximately halfway between 0 and t_1.	1 point is earned for an acceptably positioned X.
--	--

AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 3 (continued)

- (ii) Describe the change that was made to the system at time t_1 .

Additional $\text{SO}_2(g)$ was injected into the reaction vessel.	1 point is earned for the correct answer.
--	---

- (iii) After the change was made at time t_1 , the partial pressure of $\text{SO}_3(g)$ increased while the partial pressure of $\text{O}_2(g)$ decreased. Explain this observation.

The increase in the partial pressure of SO_2 perturbed the equilibrium, which responded by consuming O_2 and producing more SO_3 as new equilibrium concentrations were established. OR At equilibrium (before t_1), the reaction quotient Q is equal to K. Additional $\text{SO}_2(g)$ increases P_{SO_2}, increasing the value of Q such that $Q > K$. The reaction proceeds in reverse to reestablish equilibrium, consuming O_2 and producing more SO_3.	1 point is earned for a correct explanation.
--	--

- (e) Use the information above to answer the following.

- (i) Write the expression for the equilibrium constant, K_p , for the reaction.

$K_p = \frac{(P_{\text{SO}_2})^2(P_{\text{O}_2})}{(P_{\text{SO}_3})^2}$	1 point is earned for the correct K_p expression.
---	---

- (ii) Determine the value of K_p for the reaction.

$K_p = \frac{(0.33)^2(0.17)}{(0.50)^2} = 0.074$ OR $K_p = \frac{(0.41)^2(0.14)}{(0.57)^2} = 0.072$	1 point is earned for substituting appropriate values from the graph. 1 point is earned for correctly calculating the value of K_p.
---	--

AP[®] CHEMISTRY 2019 SCORING GUIDELINES

Question 4




The reaction between hypochlorous acid and water is represented above.

- (a) Identify one of the conjugate acid-base pairs in the reaction.

HClO and ClO [−] OR H ₂ O and H ₃ O ⁺	1 point is earned for a correct answer.
---	---

- (b) Shown in the graph below is the titration curve that results when 25.0 mL of 0.200 *M* HClO is titrated with 0.200 *M* NaOH. Carefully draw a second curve on the graph that would result from the titration of 25.0 mL of 0.200 *M* HClO₄ with 0.200 *M* NaOH.



See dashed curve above. The curve starts below the HClO curve, rises slowly from 0 to 20 mL, then steeply reaches equivalence at pH 7 and 25 mL of NaOH added.	1 point is earned for a curve that has the shape of a titration curve, that starts at a pH below the first curve, and that ends above pH 7. 1 point is earned for a curve that shows an equivalence point at 25 mL.
---	--

- (c) A student proposes creating a buffer by dissolving 0.010 mol of NaClO₄(s) in 100. mL of 0.100 *M* HClO₄. Explain why the resulting solution would not be a buffer.


A buffer solution contains a large concentration of both members in a weak acid/weak base conjugate pair. This solution would not be a buffer because HClO₄ is a strong acid, therefore ClO₄[−] is negligibly basic and cannot react with added H₃O⁺.	1 point is earned for a valid explanation.
---	---

AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 5

Half-reaction	E° (V)
$\text{O}_2 + 4 \text{H}^+ + 4 e^- \rightarrow 2 \text{H}_2\text{O}$	+0.815
$\text{NAD}^+ + \text{H}^+ + 2 e^- \rightarrow \text{NADH}$	−0.320

In human cells a compound known as NADH is involved in a reaction that generates an electrical potential. The reaction is based on the two half-reactions shown in the table above. The balanced equation for the overall reaction is shown below.



- (a) How many moles of electrons are transferred for each mole of O_2 that reacts? Justify your answer in terms of the balanced equation and the change in oxidation number of oxygen.

Four moles of electrons are needed to convert the two moles of O atoms in one mole of O_2 from the zero oxidation state to the −2 oxidation state.	1 point is earned for the correct answer.
---	---

- (b) Calculate the electrochemical potential of the overall reaction.

$E^\circ_{\text{cell}} = +0.815 \text{ V} - (-0.320 \text{ V}) = +1.135 \text{ V}$	1 point is earned for the correct answer.
--	---

- (c) Calculate the value of ΔG° for the overall reaction.

$\Delta G^\circ = -nFE^\circ$ $= -\frac{4 \text{ mol } e^-}{1 \text{ mol}_{\text{rxn}}} \times \frac{96,485 \text{ C}}{1 \text{ mol } e^-} \times \frac{1.135 \text{ J}}{1 \text{ C}}$ $= -4.380 \times 10^5 \text{ J/mol}_{\text{rxn}} = -438.0 \text{ kJ/mol}_{\text{rxn}}$	1 point is earned for the correct answer.
---	---

- (d) When the concentration of NADH increases while the concentration of NAD^+ decreases, will the electrochemical potential increase, decrease, or remain the same? Justify your answer.

Increase. The highly negative ΔG° indicates that the value of K for this reaction is very large, so decreasing Q causes the system to deviate further from equilibrium, thus increasing the magnitude of the electrochemical potential.	1 point is earned for the correct answer with a valid justification.
---	--

AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 6

Some building materials contain small capsules filled with paraffin wax to improve the insulating properties of the materials. The melting and freezing of paraffin inside the capsules in the material helps to regulate temperature. Paraffin wax begins to melt at 37°C.

- (a) While the wax is melting, is the net flow of thermal energy from the wax to the surroundings or from the surroundings to the wax? Justify your answer.

Thermal energy flows from the surroundings to the wax. Melting is an endothermic process that requires the flow of thermal energy into the material for the phase change to occur.	1 point is earned for the correct answer with justification.
--	--

- (b) Calculate the amount of thermal energy, in kJ, that is required to melt 15.2 grams of solid paraffin wax when the temperature of the surroundings is above the melting point of paraffin. (The molar mass of paraffin is 282.62 g/mol, and its molar heat of fusion is 48.78 kJ/mol.)

$15.2 \text{ g} \times \frac{1 \text{ mol}}{282.62 \text{ g}} \times \frac{48.78 \text{ kJ}}{1 \text{ mol}} = 2.62 \text{ kJ}$	1 point is earned for the correct answer.
--	---

- (c) When paraffin wax melts, its density decreases. Provide a particle-level explanation of this observation.

In liquid paraffin, the average distance between molecules is greater than in the solid state. Thus, the same number of molecules occupies a larger volume, and the density decreases.	1 point is earned for citing the greater average distance between particles. 1 point is earned for relating the increased volume to decreased density.
--	---

AP[®] CHEMISTRY
2019 SCORING GUIDELINES

Question 7


A student is given a solution of phenol red of unknown concentration. Solutions of phenol red are bright pink under basic conditions.

- (a) The student uses a pH probe and determines that the pH of the solution is 11.20. Calculate the hydrogen ion concentration, $[H^+]$, in the solution of phenol red.

$[H^+] = 10^{-pH} = 10^{-11.20} = 6.3 \times 10^{-12} M$	1 point is earned for the correct answer.
--	---

The student analyzes a sample of the solution using a spectrophotometer set at a wavelength of 559 nm, the wavelength of maximum absorbance for phenol red. The measured absorbance of the phenol red solution at 559 nm is 0.35.

- (b) Based on the calibration curve shown below, what is the concentration of the solution in micromoles per liter (μM) ?



5.4 μM (values between 5.2 μM and 5.6 μM are acceptable)	1 point is earned for a correct answer.
--	---

- (c) If the student mixed 10 mL of distilled water with 10 mL of the sample, would this diluted solution have an absorbance greater than, less than, or equal to the absorbance of the original solution? Justify your answer.

Less than. The diluted solution will have a lower concentration and therefore a lower absorbance, in accordance with Beer's law.	1 point is earned for the correct answer with a valid justification.
---	--

- (d) If the student measured the absorbance of the solution at a wavelength of 650 nm, would the absorbance be greater than, less than, or equal to the absorbance of the solution at 559 nm? Justify your answer.

Less than. 559 nm is the wavelength of maximum absorbance. The absorbance at any other wavelength will be lower.	1 point is earned for the correct answer with a valid justification.
---	--

2019 AP Chemistry Scoring Worksheet

Section I: Multiple Choice

$$\frac{\text{Number Correct}}{\text{(out of 50)}} \times 1.0000 = \frac{\text{Weighted Section I Score}}{\text{(Do not round)}}$$

Section II: Free Response

$$\text{Question 1 } \frac{\text{_____}}{\text{(out of 10)}} \times 1.0869 = \frac{\text{_____}}{\text{(Do not round)}}$$

$$\text{Question 2 } \frac{\text{_____}}{\text{(out of 10)}} \times 1.0869 = \frac{\text{_____}}{\text{(Do not round)}}$$

$$\text{Question 3 } \frac{\text{_____}}{\text{(out of 10)}} \times 1.0869 = \frac{\text{_____}}{\text{(Do not round)}}$$

$$\text{Question 4 } \frac{\text{_____}}{\text{(out of 4)}} \times 1.0869 = \frac{\text{_____}}{\text{(Do not round)}}$$

$$\text{Question 5 } \frac{\text{_____}}{\text{(out of 4)}} \times 1.0869 = \frac{\text{_____}}{\text{(Do not round)}}$$

$$\text{Question 6 } \frac{\text{_____}}{\text{(out of 4)}} \times 1.0869 = \frac{\text{_____}}{\text{(Do not round)}}$$

$$\text{Question 7 } \frac{\text{_____}}{\text{(out of 4)}} \times 1.0869 = \frac{\text{_____}}{\text{(Do not round)}}$$

$$\text{Sum} = \frac{\text{_____}}{\text{Weighted Section II Score (Do not round)}}$$

Composite Score

$$\frac{\text{Weighted Section I Score}}{\text{_____}} + \frac{\text{Weighted Section II Score}}{\text{_____}} = \frac{\text{Composite Score (Round to nearest whole number)}}{\text{_____}}$$

AP Score Conversion Chart
Chemistry

Composite Score Range	AP Score
82-100	5
70-81	4
52-69	3
37-51	2
0-36	1

2019 AP Chemistry Question Descriptors and Performance Data

Multiple-Choice Questions

Question	Skill	Learning Objective	Topic	Key	% Correct
1	1.A	ENE-4.A	Introduction to Entropy	D	89
2	4.C	SPQ-3.A	Solutions and Mixtures	B	86
3	4.A	TRA-6.B	Direction of Reversible Reactions	C	83
4	6.E	SAP-2.A	Periodic Trends	B	83
5	6.C	SAP-5.A	Intermolecular Forces	B	77
6	6.E	SPQ-3.C	Separation of Solutions and Mixtures Chromatography	D	90
7	5.F	TRA-7.D	Properties of the Equilibrium Constant	A	63
8	4.C	ENE-1.A	Catalysis	B	82
9	1.A	SAP-3.C	Structure of Ionic Solids	B	94
10	5.E	TRA-1.B	Net Ionic Equations	D	80
11	5.F	ENE-6.B	Cell Potential and Free Energy	C	80
12	5.F	SPQ-4.A	Stoichiometry	D	76
13	5.B	TRA-3.C	Concentration Changes Over Time	B	54
14	6.C	SAP-1.B	Photoelectron Spectroscopy	B	70
15	5.E	SPQ-4.A	Stoichiometry	D	90
16	6.E	ENE-3.A	Bond Enthalpies	D	73
17	6.E	SAP-5.A	Intermolecular Forces	D	76
18	5.F	ENE-2.F	Introduction to Enthalpy of Reaction	C	50
19	4.A	SAP-7.B	Kinetic Molecular Theory	B	72
20	6.D	ENE-4.C	Gibbs Free-Energy and Thermodynamic Favorability	D	67
21	6.E	SAP-10.B	Properties of Buffers	B	74
22	5.E	TRA-1.C	Representations of Reactions	B	86
23	5.F	TRA-3.C	Concentration Changes Over Time	A	67
24	5.D	SAP-3.B	Intramolecular Force and Potential Energy	B	48
25	5.C	SAP-7.A	Ideal Gas Law	A	68
26	6.E	SAP-5.A	Intermolecular Forces	A	55
27	4.A	SAP-9.C	Weak Acid and Base Equilibria	B	60
28	4.A	SPQ-4.B	Introduction to Titration	C	31
29	5.F	SAP-9.A	Introduction to Acids and Bases	B	33
30	4.A	ENE-2.C	Heat Transfer and Thermal Equilibrium	D	67
31	4.C	SAP-2.B	Valence Electrons and ionic Compounds	D	57
32	5.F	SAP-9.A	Introduction to Acids and Bases	C	30
33	5.B	ENE-4.C	Gibbs Free-Energy and Thermodynamic Favorability	D	36
34	6.F	SPQ-5.A	Introduction to Solubility Equilibria	D	48
35	5.E	TRA-1.B	Net Ionic Equations	B	55
36	5.F	SPQ-2.B	Composition of Mixtures	D	42
37	5.F	SPQ-3.A	Solutions and Mixtures	D	25
38	1.B	TRA-3.A	Reaction Rates	D	57
39	5.E	SAP-9.C	Weak Acid and Base Equilibria	C	64

2019 AP Chemistry Question Descriptors and Performance Data

Question	Skill	Learning Objective	Topic	Key	% Correct
40	6.E	ENE-4.D	Thermodynamic and Kinetic Control	C	65
41	4.A	SAP-9.E	Acid-Base Titrations	B	47
42	1.A	SAP-4.C	VSEPR and Bond Hybridization	C	52
43	1.A	SAP-4.C	VSEPR and Bond Hybridization	B	28
44	1.A	SAP-5.B	Properties of Solids	A	51
45	1.A	SAP-3.A	Types of Chemical Bonds	C	45
46	2.C	ENE-2.D	Heat Capacity and Calorimetry	C	57
47	5.F	TRA-8.B	Reaction Quotient and Le Châtelier's Principle	A	68
48	6.E	SAP-5.A	Intermolecular Forces	D	41
49	5.F	SPQ-5.B	Common-Ion Effect	B	34
50	5.F	TRA-7.D	Properties of the Equilibrium Constant	D	56

Free-Response Questions

Question	Skill	Learning Objective	Topic	Mean Score
1	1.A 1.B 4.A 5.F 6.E	TRA-2.A SAP-7.A SPQ-4.A	4.7 3.4 4.5 6.4 6.6 3.1 3.77.9	5.98
2	3.B 5.D 3.A 5.F 6.F 6.G 5.E	TRA-1.C ENE-2.B TRA-4.C	4.3 6.2 5.6 1.1 4.2 5.3	5.62
3	1.A 2.D 3.B 3.C 6.D 5.B 5.F	SAP-4.A SAP-4.C ENE-4.C TRA-6.A TRA-7.A TRA-7.B TRA-8.A	2.5 2.7 9.3 7.1 7.3 7.4 7.9	5.97
4	1.A 3.A 4.A	TRA-2.B SAP-9.E SAP-10.B	4.8 8.5 8.8	2.22
5	4.A 5.F	TRA-2.A ENE-6.B ENE-6.C	4.7 9.8 9.9	1.79
6	4.A 4.C 5.F	ENE-2.A ENE-2.E SAP-5.B	6.1 6.5 3.2	2.07
7	5.F 2.B 2.D	SAP-9.A SAP-8.C	8.1 3.13	2.28